



ISSN: 2347-6567

International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)

IJAMSCR | Vol.11 | Issue 4 | Oct - Dec -2023

www.ijamscr.com

DOI : <https://doi.org/10.61096/ijamscr.v11.iss4.2023.511-522>

Research

Antidiabetic Activity of Methanolic Extracts of Leaves of *Tylophora neglecta* Leaves in Alloxan Induced Diabetes in Rats

*Mohammed Abdul Rahman Khalid, Mohammed Abdul Aziz Shahid, Sollar Murali

*KGR Institute of Technology and Management, Rampally, Affiliated to Osmania University, Hyderabad, Telangana-50130, India.

*Correspondent author: Mohammed Abdul Rahman Khalid

Email: crazy4ukhalid@gmail.com

	Abstract
Published on: 1 Dec 2023	Diabetes mellitus is an endocrine disorder that negatively influences about 300 million humans worldwide. Pharmacological activity carried out on <i>Tylophora neglecta</i>'s for an antidiabetic activity in Alloxan-induced diabetes in albino mice used to be evaluated with grouping the animals into six organizations (n=6). Intraperitoneal implantation occurred sixteen hours prior to the onset of diabetes. After extensive health testing, alloxan used to set off experimental diabetes in Swiss mice and blood sugar degrees before and after administration of normal, untreated, and diabetic mice. Quantitative techniques of records series and analysis allowed for a thorough evaluation. No detrimental results were considered at oral doses of 2000mg/kg physique weight in the brilliant oral threat reviews of the concentrates. The hypoglycemic and antidiabetic results of <i>Tylophora neglecta</i> aqueous and alcoholic concentrates of 20 and 30mg/kg body weight have been tested. In this study, 72 hours of fasting preceded via a single Intraperitoneal (IP) dose of Alloxan monohydrate resulted in diabetes in animals. The FBGL used to be measured on days 7, 14, and 21, and the OGTT was once performed on days 8, 15, and 22. The outcomes with glibenclamide are quite comparable to the norm. Research has showed that the leaves of <i>Tylophora neglecta</i> have antidiabetic recreation by using merchandising the regeneration of cells accountable for secreting the hormone insulin.
Published by: DrSriram Publications	
2023 All rights reserved.	
	
Creative Commons Attribution 4.0 International License.	Keywords: Tylophora neglecta, Alloxan, Antidiabetic, Total antioxidant

INTRODUCTION

Diabetes is one of the most common non-communicable diseases and a serious life-long condition appearing worldwide. The etiology of diabetes is a complex interaction of genetic and environmental factors. It is a heterogeneous group of metabolic disorders characterized physiologically by dysfunction of pancreatic beta cells and deficiency in insulin secretion or insulin activity and clinically by hyperglycemia or impaired glucose tolerance and other manifestable disorders. It is an endocrinological syndrome abnormally having high levels of sugar in the blood. This may be either due to insulin not being produced at all, is not made at sufficient levels, or is not as effective as it should be Diabetes is still a serious health problem all over the world since it is associated

with increased morbidity and mortality rate. When compared with the general population, mortality and morbidity increase in diabetes is mainly due to the associated chronic complications both specific (micro vascular) and nonspecific (macro vascular). Since the disease prevails in both genders and in all age groups, the general public has a concern about its control and treatment.

PLANT PROFILE

Botanical Name: *Tylophora neglecta*

Kingdom: Plantae

Order: Gentianales

Family: Apocynaceae

Sub family: Asclepiadaceae

Genus: Tylophora

Uses: People take *Tylophora* by mouth for allergies, asthma, cancer, congestion, constipation, cough, inflamed skin, diarrhea, bloody diarrhea, gas, hemorrhoids, tender joints (gout), yellowed skin (jaundice), joint disorder (rheumatoid arthritis), whooping cough, to make someone vomit, and to cause sweating.

AIM AND OBJECTIVES

Evaluation of the alcoholic concentrates of *Tylophora neglecta* leaves for their Phytochemical and pharmacological properties. An effort, ongoing during the work, was once completed to evaluate the unpleasantness of diabetic activity. *Tylophora neglecta* water and alcohol concentrations are being planned. To simulate the phytochemical comparison of the two extracts, to use these concentrations in a wildly unfavorable manner. To hint the improvement of diabetes-related resistance in comparable rodents. Alloxan-induced diabetic rats were chosen for the counter-diabetic enhancement study.

MATERIALS AND METHODS

Experimental animals: Wistar rats of both sexes were selected for learn about due to the fact they are large, healthy, and without problems handled. The animals had been fed a pellet meal (the Amrul Exploration office's encouraged food) and housed in clean, secure locations with ample fencing and no get admission water. They have been avoided for the time being earlier than to the day of evaluation, following a speedy of 72 hours beginning on the day of the Alloxan show. The creature domestic used to be maintained at a consistent 27 tiers Celsius to make sure the alleviation of the animals housed within. IAEC assurances have been present in animal testing.

Plant Material Collection: After being recognized and tested the usage of the plant's Part of Pharmacognosy, the leaves of *Tylophora neglecta* have been gathered at a local market in Hyderabad at some point of the month of January. After being cleansed, the plant remember was reduce into smaller pieces, let to air dry at room temperature whilst covered, and then blended into a coarse powder. Hiding the powdered substance till it may want to be extracted from its hiding place.

Preparation of plant extracts

Preparation of Aqueous Extract: About 20 gm of dried *Tylophora neglecta* leaves have been introduced to a 250ml receptacle that contained about 200ml of water. After the elements were mixed, they have been let to upward push for 4 to 5 hours at eighty to 900 tiers Celsius. Whatmann channel paper was once then used to in addition separate the concentrate. The filtrate used to be focused till the step forward should be shaped. The centered substance was once sealed in pattern containers, kept at room temperature, and sooner or later used for further evaluation of the exercises.

Preparation of Alcoholic Extract: About 20g of dried *Tylophora neglecta* leaves had been added to a 250-milliliter container conserving 200 ml of liquor. The components had been mixed thoroughly, then let to upward jostle for 4 to five hours at temperatures between fifty and six hundred stages Celsius. Whatmann channel paper was then used to further separate the concentrate. The filtrate was targeted until the step forward should be shaped. Prior to being used for similarly investigation into the activities, the targeted item was once saved in mannequin covers at room temperature.

Preliminary phytochemical analysis of the extracts: The resulting concentrates were fed into a phytochemical screening system. The following phytochemical analyses have been performed to analysis phytoconstituents:

Alkaloids :(a. Mayer's test, b. Wagner's test , c. Dragendroff's tested. Hager's test), Triterpenoids: (Salkowski's Test), Saponins (Froth Test), b. Liberman Burchard Test, Flavonoids (Alkaline reagent test, Lead acetate test), Phenolic and Tannins (Ferric chloride test, Gelatin test, Vanillin hydrochloride Test)

Selection of dose for animal study: *Tylophora neglecta* (3-5 g/kg) physique phase used to be sourced from a human donor for the reason of rodent evaluation. According to Ghosh (1984), a rat has a 0.018 human piece variant (per 2 hundred g physique weight). Rat doses of 20 and 30 mg/kg have been decided after comparing them to the human doses of 0.3 and 0.5 g/kg. At 2000 milligrams per kilogram of physique weight, extreme toxicity was eliminated.

Pharmacological evaluation

Preparation of extracts: Alcoholic and non-alcoholic concentrations of the plant *Tylophora neglecta*, suspended in water with 3% v/v Tween-80. For the sake of clarity, all of the were managed orally. When necessary, plans for the concentrations were synchronized. Ten milliliters per kilogram was once chosen as the well-known for all animals.

Acute Oral Toxicity: *Tylophora* fluid alcoholic concentrates' severe oral toxicity was decided the use of Wistar rats (200-250g) that had been well cleansed and stored in regular conditions. There have been 425 human beings employed to head up harm investigations. The animals have been given a single dose of man or woman listen (up to 2000 mg/kg), and their mortality all through the next two and seven days was recorded. Over the path of a week, they additionally had direct and neurological profiles examined.

Assessment of Anti-diabetic Activity in Normal and Alloxan induced Rats: The animals, which have been arbitrarily divided into six groups of four, all refused to improve at a faster clip. At zero hours, earlier than the I.P. injection of concentrates/standard/vehicle, blood samples have been drawn from the patient's tail vein for analysis. Then, on days zero, seven, fourteen, and twenty-one, blood was once drawn at intervals of 2, 4, and 8 hours after the meal and analyzed the usage of a glucometer (ONE TOUCH glucometer) to determine blood glucose levels.

Oral glucose tolerance test (OGTT) in normal rats: One hour after the euthanization/Glibenclamide/vehicle association, all animals in all businesses had been given 2g/kg of glucose. Blood samples have been taken from the tail vein at zero hours, 1/2 hours, 1 hour, 1 half of hours, and two hours after the glucose load was once administered. Individual glucometer measurements had been recorded on days 1, 8, 15, and 22 to decide blood glucose levels.

Assessment of Anti-Diabetic Activity in Alloxan Induced Diabetic Rats: Both sexes of the rats with mild fur and a weight between 200 and 250 g had been covered in the inspection. All of the animals have been housed in rat cages at room temperature and fed pellets and water except restriction, Alloxan used to be extracted from everyday saline. Using an Intraperitoneal injection of one hundred twenty mg/kg of alloxan in everyday saline, we brought about diabetes in rats that fasted for sixteen hours. After alloxanization, the rats had been administered a 5% w/v glucose administration sketch in their cages for the following 24 hours to stop hypoglycemia. Research the usage of rats with fasting blood glucose levels above 2 hundred mg/dl have been selected after 72 hours. These animals were chosen after being monitored for seven days to ensure they had strong hyperglycemia (fasting blood glucose ranges higher than two hundred mg/dl and under four hundred mg/dl). They had been then divided into six companies of 4 to be used in the resultant research on the preliminary models. Each of the aforementioned companies of animals adhered to the days cited treatment norm. At 0, 1, 2, 4, and eight hours after the affiliating, blood samples were taken via retro orbital incision. For the next 22 days, therapy was once administered. Additional blood samples were collected on days 7, 14, and 21 after the 1-hour relationship for secondary analysis. The glucometer used to be used at a number of intervals to decide the blood sugar levels.

Oral glucose tolerance test (OGTT) in alloxan induced diabetic rats: Similar Alloxan-injured diabetic animals have been subjected to OGTT on days 8, 15, and 22 for contrast investigations of diabetes progression.

In every group of animals, two g/kg of glucose was once administered an hour after euthanasia, Glibenclamide administration, or car transport. Blood samples have been taken thru retro orbital slice at 0, 0.5, 1, and two hours after the glucose load was administered. Blood samples had been taken from the animal's tail vein, and its glucose ranges have been measured with a glucometer device.

STATISTICAL ANALYSIS

One-way evaluation of variance (ANOVA) and the T-test have been used to analyze the data, which used to be as capability and widespread blunders of the means. There were two one-of-a-kind examination plans in place. For instance, compare one dealt with bundle to a manipulate group. Second, the Diabetic Supervision team vs every

handled pack. Significant differences between the organizations had been regarded to exist at the P0.001 and P0.05 levels, respectively.

RESULTS

Phytochemical screening of *Tylophora neglecta*: The persevering with investigation has assumed that the remotod mixes from the plant *Tylophora neglecta* existing diverse pharmacological open problems due to the existence of a range of phytochemical components. Constraints imposed by means of the plant's substances necessitate extra study, and any pharmacological enhancements made to the plant should be done so for the benefit of humankind.

Table1: Phytochemical screening of *Tylophoraneglecta*.

S.No.	Phytoconstituents	Aqueous	Alcoholic
1.	Alkaloids	-	-
2.	Flavonoids	+	+
3.	Steroids	+	-
4.	Tannins	+	+
5.	Anthraquinones	-	-
6.	Terpenoids	+	+
7.	Cardiac glycoside	+	+
8.	Saponins	+	-

Acute toxicity testing: *Tylophora neglecta* ethanol concentrations were shown to be secure up to 2000 mg/kg body weight, and their LD 50 was calculated to be higher than 2000 mg/kg. Up to the very quiet of the time duration beneath study, no fatalities or hazardous reactions have been observed.

HYPOGLYCEMIC ACTIVITY IN NORMAL RATS: On day 0 of the study, the FBGL for each crew was once in vary of ninety to one zero five mg/dl. Concentrate exhibits critical hypoglycemic motion with repeated association, as proven by means of the great discount in blood glucose stages after repeated therapy on days 7, 14, and 21 with dosages of watery and alcoholic concentrate (100 and 2 hundred mg/kg). Table No. 10 summarizes the FBGL shifts throughout activities following the reorganization of sections. The FBGL used to be drastically (p0.005) lower on days 7, 15, and 21 after repeated affiliation with each watery and alcoholic concentrates, indicating that the latter can transfer hypoglycemia on recurrent affiliation. However, with Glibenclamide as a distinction and other groups, hypoglycemia growth was most suggested on days 7, 14, and 21. The consequences of the repeated section association advocate that the liquid and alcoholic concentrates have good sized hypoglycemic growth.

Effect of extracts of *Tylophora neglecta* on fasting blood glucose level (FBGL) in normal rats

Table 2: Impact of concentrates of *Tylophora neglecta* on fasting blood glucose level (FBGL) in typical rodents.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		7 th day	14 th day	21 st day
Normal control	-	87.15±1.16	75.33±2.98	64.15±1.27
Glibenclamide	10	81.89±3.73	76.82±3.26	66.16±2.09
AQTN1	20	85.30±1.92	80.60±1.64	71.18±5.22
AQTN2	30	80.78±3.47	72.85±5.81	67.59±2.94
ALTN1	20	76.91±1.20	61.27±1.68	55.17±5.79
ALTN2	30	82.5±1.29	70.43±2.21	65.23±1.58




Fig 1: Effect of extracts of *Tylophoraneglecta* on fasting blood glucose level (FBGL) in normal rats.

Table 3: On the seventh day, the effects of concentrates of *Tylophoranelecta* on normal rats' fasting blood glucose level (FBGL).

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl) (7 th day)			
		1 st hr	2 nd hr	4 th hr	8 th hr
Normal control	-	40.16±2.08	52.71±1.25	76.14±0.89	87.13±1.14
Glibenclamide	10	50.14±1.39	61.21±1.20	70.2±0.05	81.87±3.78
AQTN1	20	53.52±0.76	65.31±1.03	72.54±8.52	85.31±1.91
AQTN2	30	47.72±1.25	60.59±2.87	71.33±1.06	80.76±3.49
ALTN1	20	45.1±0.54	58.35±1.50	64.25±3.28	76.92±1.19
ALTN2	30	53.1±0.08	65.28±6.19	76.10±2.75	82.1±1.24




Fig 2: The influence of *Tylophoranelecta* concentrates on the fasting blood glucose level (FBGL) of ordinary rats on the seventh day of the experiment.

Table 4: Impact of concentrates of *Tylophora neglecta* on fasting blood glucose level (FBGL) in ordinary rats on fourteenth day

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl) (14 th day)			
		1 st hr	2 nd hr	4 th hr	8 th hr
Normal control	-	44.15±2.91	57.19±0.14	74.15±2.16	75.32±2.97
Glibenclamide	10	52.05±1.56	65.25±2.25	69.26±1.30	75.81±3.25
AQTN1	20	48.19±3.34	62.63±1.01	75.55±2.03	80.61±1.61
AQTN2	30	55.36±1.78	57.10±3.51	69.93±2.16	72.86±5.84
ALTN1	20	50.53±2.81	55.02±1.41	61.30±0.28	61.23±1.69
ALTN2	30	51.21±3.67	66.98±2.02	68.45±3.12	70.46±2.26




Fig 3: Effect of extracts of *Tylophoranelecta* on fasting blood glucose level (FBGL) in normal rats on 14th day

Table 5: Effect of extracts of *Tylophoranelecta* on fasting blood glucose level (FBGL) in normal rats on 21st day

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl) (21 st day)			
		1 st hr	2 nd hr	4 th hr	8 th hr
Normal control	-	38.45±1.22	43.03±1.70	56.14±2.95	64.15±1.27
Glibenclamide	10	41.27±2.03	54.26±2.08	58.08±1.23	66.07±2.06
AQTN1	20	36.15±0.58	43.11±1.02	52.13±3.07	71.18±5.21
AQTN2	30	42.11±3.13	51.04±2.08	64.31±2.36	67.57±2.94

ALTN1	20	34.03±1.28	40.32±2.37	53.42±0.22	55.14±5.72
ALTN2	30	39.17±4.04	44.23±1.12	51.21±3.04	65.20±1.56




Fig 4: Effect of extracts of *Tylophora neglecta* on fasting blood glucose level (FBGL) in normal rats on 21st day

Oral glucose tolerance test (OGTT): Liquid and alcoholic concentrates of *Tylophora neglecta* usually (P0.005) suppress the rising in FBGL after glucose load (2g/kg) in rodents for as lengthy as two hours, compared to other agencies that take away Glibenclamide on days 8, 15, and 22. While both aqueous and alcoholic concentrates notably diminished FBGL levels. Large (P0.005) increases in FBGL were viewed 30 minutes, 1 hour, and two hours after a single dose of glibenclamide (10mg/kg).

Table 6: Effect of extracts of *Tylophoraneglecta* on 8th, 15th and 22nd day in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		8 th day	15 th day	22 nd day
Normal control	-	85.06±3.73	90.37±1.85	92.32±2.94
Glibenclamide	10	80.12±2.78	75.18±2.73	70.22±1.51
AQTN1	20	86.27±1.22	81.01±0.39	76.18±4.76
AQTN2	30	78.68±0.34	73.98±2.55	71.56±2.27
ALTN1	20	82.56±1.75	78.26±1.92	70.39±3.75
ALTN2	30	71.29±2.57	66.47±1.42	60.62±1.67




Fig 5: Effect of extracts of *Tylophora neglecta* on 8th, 15th and 22nd day in normal rats.

Table 7: Effect of extracts of *Tylophora neglecta* on 8th day in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)8 th day			
		0.5 hour	1 st hour	1.5 hour	2 nd hour
Normal control	-	36.26±2.61	63.15±1.12	71.01±2.20	85.05±3.72
Glibenclamide	10	32.10±0.52	60.07±2.26	67.22±3.19	80.11±2.79
AQTN1	20	42.32±1.02	53.25±1.15	80.03±2.54	86.26±1.21
AQTN2	30	50.15±2.53	58.08±0.10	67.15±2.19	78.67±0.33
ALTN1	20	56.20±1.55	62.19±2.30	78.31±1.82	82.57±1.76
ALTN2	30	50.12±1.63	59.28±1.00	65.24±5.02	71.28±2.58




Fig 6: Effect of extracts of *Tylophoranelectaon* 8th day in normal rats.

Table 8: Effect of extracts of *Tylophoranelectaon* 15th day in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl) 15 th day			
		0.5 hour	1 st hour	1.5 hour	2 nd hour
Normal control	-	41.11±3.21	62.12±1.19	81.25±1.19	90.36±1.84
Glibenclamide	10	35.06±0.12	60.26±2.12	76.11±2.25	75.17±2.74
AQTN1	20	46.15±2.10	58.33±0.21	72.01±0.01	81.02±0.36
AQTN2	30	53.60±1.15	61.18±2.21	65.19±1.25	73.99±2.54
ALTN1	20	62.22±3.00	68.02±1.00	73.20±0.11	78.25±1.91
ALTN2	30	45.15±2.23	56.28±1.25	62.38±0.32	66.46±1.44




Fig 7: Effect of extracts of *Tylophora neglecta* on 15th day in normal rats

Table 9: Effect of extracts of *Tylophora neglecta* on 22nd day in normal rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl) 22 nd day			
		0.5 hour	1 st hour	1.5 hour	2 nd hour
Normal control	-	47.18±1.24	69.1±2.11	87.24±2.90	92.31±2.93
Glibenclamide	10	41.38±2.18	67.28±1.10	70.28±5.17	70.21±1.50
AQTN1	20	51.19±1.28	65.34±1.01	78.54±1.42	76.17±4.75
AQTN2	30	58.01±3.12	63.22±3.56	68.39±2.14	71.57±2.26
ALTN1	20	60.58±1.35	65.26±2.14	72.15±0.78	70.38±3.76
ALTN2	30	38.15±3.57	46.25±1.35	52.23±1.28	60.61±1.65




Fig 8: Effect of extracts of *Tylophora neglecta* on 22nd day in normal rats

Anti-diabetic activity in alloxan induced diabetic rats: Most rats had fasting blood glucose levels (FBGL) between ninety and one hundred mg/dl. After 72 hours, the FBGL had elevated to 252-266 mg/dl thanks to alloxan therapy (120 mg/kg, I.P.). On day seven, these characteristics had been normalized to a degree of 255 mg/dl. A table depicts versions in fasting blood glucose degrees across a range of groups. This records proven that the blood glucose levels of the normal manage critters had been secure throughout the period of the study. The fasting blood glucose stages of the diabetic reference group had improved dramatically over the past 21 days. Fasting blood glucose degrees have been substantially decreased ($p < 0.05$) in the glibenclamide (10mg/kg) treatment team on days 7, 14, and 21 of the research.

Effect of *Tylophora neglecta* extracts on antidiabetic activity in Alloxan induced diabetic rats: On days 7, 14, and 21, animals from separate agencies treated with one hundred and 200mg/kg of fluid and alcoholic of more than a few concentrations showed a vast reduction (P0.05) in FBGL. When in contrast to alcohol concentrates at the same social event, watery concentrates have dropped more (percentage wise) in FBGL.

Table 10: Effect of extracts of *Tylophora neglecta* on fasting blood glucose level (FBGL) in Alloxan induced diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		7 th day	14 th day	21 st day
Normal control	-	81.18±4.67	80.91±7.88	82.75±4.36
Diabetic control	10	295.11±11.06	268.05±10.04	251.14±12.56
Glibenclamide	10	262.22±10.17	245.19±12.02	230.27±11.84
AQTN1	20	373.63±11.22	361.17±19.63	332.56±29.12
AQTN2	30	381.24±12.35	368.84±17.17	351.99±38.34
ALTN1	20	292.09±13.08	275.04±14.29	222.61±05.92
ALTN2	30	208.44±12.18	189.28±16.17	160.31±20.35




Fig 9: Effect of extracts of *Tylophora neglecta* on fasting blood glucose level (FBGL) in Alloxan induced diabetic rats.

Oral glucose tolerance test (OGTT) on 8th, 15th and 22nd day-The increase in FBGL following a glucose load (2g/kg) in mice is generally (P0.05) blanketed via the fluid and alcoholic concentrates of *Tylophora neglecta* in the first 30 minutes and for up to 2 hours as distinction and one-of-a-kind businesses discard Glibenclamide on days 8, 15, and 22. While each alcoholic and water-based concentrates considerably decreased FBGL, Glibenclamide (10 mg/kg) demonstrated a vast (P 0.05) reduction in FBGL upward push at 30 minutes, 60 minutes, and 2 hours.

Table 11: Effect of extracts of *Tylophora neglecta* on 8th, 15th and 22nd day in Diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)		
		8 th day	15 th day	22 nd day
Normal control	-	87.75±3.63	84.59±2.87	90.99±1.58
Diabetic control	10	285.01±02.69	275.00±19.55	321.57±14.99
Glibenclamide	10	362.38±21.10	312.88±11.73	289.41±18.55
AQTN1	20	260.26±10.62	241.70±16.40	208.36±11.90
AQTN2	30	287.14±13.30	280.78±13.78	255.98±19.75
ALTN1	20	261.22±12.11	221.56±17.87	188.63±17.99
ALTN2	30	353.55±14.61	330.76±15.28	270.66±10.86





Fig 10: Effect of extracts of *Tylophora neglecta* on 8th, 15th and 22nd day in Diabetic rats.

Table 12: Effect of extracts of *Tylophora neglecta* on 8th day in Diabetic rats.


Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)8 th day			
		0.5 hour	1 st hour	1.5 hour	2 nd hour
Normal control	-	51.69±2.68	63.26±2.18	75.19±2.91	87.76±3.64
Diabetic control	10	195.26±15.73	212.17±18.58	245.14±10.12	285.02±02.68
Glibenclamide	10	184.02±21.54	248.19±21.06	301.11±15.27	362.39±21.11
AQTN1	20	182.33±31.27	190.27±24.82	221.26±10.64	260.25±10.63
AQTN2	30	170.17±21.52	182.17±54.44	250.86±21.37	287.13±13.31
ALTN1	20	162.04±14.12	167.67±02.06	221.57±05.18	261.21±12.10
ALTN2	30	153.23±21.54	211.73±10.34	280.37±13.74	353.56±14.62

Table 13: Effect of extracts of *Tylophora neglecta* on 15th day in Diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl)15 th day			
		0.5 hour	1 st hour	1.5 hour	2 nd hour
Normal control	-	48.26±3.72	60.35±3.18	76.24±1.53	84.58±2.83
Diabetic control	10	186.37±15.73	201.17±12.80	231.24±12.65	275.01±19.56
Glibenclamide	10	193.45±1.32	272.15±16.13	313.29±11.32	312.89±11.74
AQTN1	20	160.12±61.11	206.51±04.88	235.33±15.74	241.71±16.42
AQTN2	30	183.76±13.63	196.26±63.11	272.03±01.18	280.79±13.77
ALTN1	20	150.14±10.76	186.08±1.14	230.16±1.22	221.55±17.84
ALTN2	30	182.13±51.97	256.76±15.02	290.12±36.14	330.77±15.26


Fig 11: Effect of extracts of *Tylophora neglecta* on 15th day in Diabetic rats.
Table 14: Effect of extracts of *Tylophora neglecta* on 22nd day in Diabetic rats.

Treatment	Dose (mg/kg)	Blood glucose level(mg/dl) 22 nd day			
		0.5 hour	1 st hour	1.5 hour	2 nd hour
Normal control	-	61.26±0.52	66.4±2.06	83.53±3.97	90.98±1.59
Diabetic control	10	203.16±02.37	232.26±50.19	281.17±33.26	321.55±14.98
Glibenclamide	10	163.88±3.45	212.73±23.04	253.18±13.07	289.42±18.56
AQTN1	20	143.42±53.08	169.23±26.18	183.57±53.38	208.37±11.91
AQTN2	30	159.58±08.19	199.18±39.03	263.12±06.18	255.99±19.76
ALTN1	20	141.12±14.08	162.88±3.53	180.74±2.64	188.64±17.98
ALTN2	30	154.18±08.12	182.15±02.02	225.64±14.26	270.67±10.88


Fig 12: Effect of extracts of *Tylophora neglecta* on 22nd day in Diabetic rats

DISCUSSION

Despite diabetes' awesome prevalence, severity, and fatality price throughout the world, the disease is frequently viewed untreatable at best. The poor consequences it has on the body can be mitigated in section through dietary adjustments, plant treatments, and a range of synthetic tranquillizers. A choose crew of authors have investigated the practicable of useful plants as anti-diabetic experts. Many plant life have been tested in-vivo using animal models, such as rodents, towards the complexities wrought with the aid of diabetes inducers, and it has been unfold that many flowers has the attainable to reduce fasting blood glucose tiers and assist in working on other diabetic captures. The threat of growing secondary diabetic complications is decreased if hyperglycemia continues to decrease. Consistent glucose administration is the key to keeping off or resolving the difficulties diabetics face and achieving non-public success.

Numerous robust combinations characteristic of plant life have been identified. Alkaloids, flavonoids, steroids, tannins, anthraquinones, terpenoids, coronary heart glycosides, and other bioactive compounds structure this complicated net of excessive standards. Against hyperglycemic and glucose suppressive activities, these combos have been proven to solid regions for convey. These effects may also be completed by using modulating insulin secretion from pancreatic β -cells, blocking the absorption of glucose with the aid of the stomach, stimulating glycogenesis in the liver, and growing the body's glucose usage. In addition to restoring enzymatic functions, repairing and recovering pancreatic islets, and dealing with hepatic and kidney damage, these combos may also show off calming, antibacterial, antifungal, and cardio cautious activities.

At concentrations of 20 and 30 mg/kg, the harsh liquid and alcoholic concentrates of *Tylophora neglecta* leaves altered glucose blockage and lowered fasting blood glucose degrees in normoglycemic mice, revealing the hypoglycemic idea of the concentrates. The impact was once greater conveyed between the two concentrations. These findings propose that hypoglycemia might also be prompted by means of the concentrates through a mechanism distinct from insulin secretion, such as the inhibition of endogenous glucose production or glucose processing.

Alloxan monohydrate is one of the engineered specialists used to trigger diabetic mellitus in animals. Damage to the Langerhans islet beta cells is a main cause of diabetes. An oxygen generator that sweeps DNA with free oxygen. A single intravenous dose of Alloxan validated baseline hyperglycemia. The underlying systemic hyperglycemia in the diabetic situation is due to excessive hepatic glycogenolysis and gluconeogenesis, which is associated with lowered utilisation of glucose via tissues. Since alloxan-induced hyperglycemia is considered moderate diabetes and can also reverse after roughly a month of therapy, we monitored the hypoglycemic impact of the plant on hyperglycemic animals for a complete month. When compared to the consequences of the gold popular drug Glibenclamide, the concentrates were viewed as having a monster effect.

Tylophora neglecta leaf extracts were phytochemically analyzed, and the consequences showed the existence of antidiabetic associate metabolites. Concentrates of this plant had the identical flavonoids, alkaloids, and steroids that gave different plant life their anti-diabetic properties. It's possible that the plant's anti-diabetic effects are due to the phenols it contains. Evidence suggests that phenols can shield against diabetes development with the aid of preventing the dying of beta cells through interfering with the peroxidation chain reaction. *Tylophora neglecta* extracts show up to be intriguing starting points for in addition research that might lead to advances in diabetes treatment. Because of their lower cost and shorter development period, Phytomedicine are a better match for our present day economic local weather than allopathic drug research and development.

SUMMARY

This ongoing research aimed to higher apprehend the antidiabetic effects of an ethanol extract of *Tylophora neglecta* leaves in streptozotocin-induced diabetic rats. The leaves of *Tylophora neglecta* L. had been organized for Phytochemical screening for Carbohydrates, alkaloids, flavonoids, glycosides, Saponins, steroids, tannins, and terpenes. The *Tylophora neglecta* leaves used in this mission had been purchased domestically after drying. *Tylophora neglecta* leaves were continually flushed down the rest room with a combination of water and alcohol. Extricates had been sought for their opposition to the improvement of diabetes in both ordinary and Alloxan-induced diabetic mice, which led to the successful decision of the restorative component of the concentrates. Limits on the journey have been assessed as follows: Normal and alloxan-induced animals' fasting blood sugar stages have been measured on days 7, 14, and 21. Glucose tolerance check (oral): In each non-stimulated and alloxan-stimulated rats, these phases happen on days 8, 15, and 22. Six, oral glucose tolerance take a look at (OGTT) in existing 2nd (day 7) and large length (day 14 and 21) rehashed relationship in everyday and alloxan induced diabetic rodents confirmed that fluid and alcoholic concentrates of *Tylophora neglecta* (20 and 30 mg/kg) had a massive have an effect on in blood glucose reducing improvement. Previous lookup has located that *Tylophora neglecta*'s aqueous and alcoholic concentrates had tremendous anti-diabetic consequences after repeated use.

CONCLUSION

The study was performed to find out the beneficial effects of two different extracts of leaves of *Tylophora neglecta* in normoglycemic rats and alloxan induced diabetic rats therefore, the present work was conceived to evaluate the phytochemical and pharmacological screening of leaves of *Tylophora neglecta*. The phytochemical evaluation has revealed the presence of Alkaloids, Flavonoids, Steroids, Tannins, Anthraquinones, Terpenoids, Cardiac glycoside.

The aqueous and alcoholic extracts had hypoglycemic activity because of the presence of flavonoids which are rich in treatment of hypoglycemia with less side effects. Flavonoids might be producing hypoglycaemic effect by a mechanism independent from insulin secretion e.g. by the inhibition of endogenous glucose production or by the inhibition of intestinal glucose absorption. The present study *Tylophora neglecta* of both aqueous and alcoholic extracts was shown significant effect on glucose tolerance and also showed reduction in fasting blood glucose levels in normal diabetic rats.

The data of the blood glucose level of rats treated with Alloxan (150mg/kg body weight) produced diabetes within 72 hours. After 72 hours of Alloxan administered the blood glucose levels of rats were observed. It was observed that significant lowering of sugar in aqueous and alcoholic extract. The administration of different extracts at a dose of 20 and 30 mg/kg showed significant anti-hyperglycemic effect at 22nd days which was evident from the 7th day onwards as compared to standard. The aqueous and alcoholic extract of *Tylophora neglecta* has shown better anti-hyperglycemic effect of the extract on the fasting blood sugar levels on diabetic rats as shown in table. The decreasing blood glucose levels are comparable with that of 10 mg/kg of Glibenclamide. The Glibenclamide (10 mg/kg body weight) shows significant effect on compared to the initial and more significant effect on the 22nd Day compared to the initial. The aqueous and alcoholic extracts of 20 and 30mg/kg body weight shows significant.

Results of anti-diabetic activity in normal and Alloxan induced rats the extract established the scientific basis for the utility of these plants in the treatment of diabetes. The extracts have shown significant reduction in blood glucose levels in normal and Alloxan induced diabetic rats and produced maximum anti-diabetic activity and are higher than the hypoglycaemic activity of Glibenclamide in the diabetic rats. In glucose loaded animals, the drug has reduced the blood glucose to the normal levels. It is possible that the drug may be acting by potentiating the pancreatic secretion or increasing the glucose uptake. In conclusion, these extracts showed significant anti-diabetic effect in normal and diabetic rats after administration. Thus the claim made by the traditional Indian systems of medicine regarding the use of these plants in the treatment of diabetes stands confirmed.

Future scope: The current thesis revealed the antidiabetic potential of *Tylophora neglecta* being effective in hyperglycemia and their ability to protect against other metabolic aberrations caused by diabetes validates its therapeutic traditional use. The antidiabetic effect of *Tylophora neglecta* may be due to their ability to increase the number and restore the integrity of pancreatic β -cells of islets of Langerhans. However further studies are required to evaluate other chemical constituents of the plant for their hypoglycemic potential and to identify their targets, mechanism of action and combination of plant products with synthetic drugs.

REFERENCES

1. Consultation, W.H.O. DefinitionDiagnClassif DiabetesMellitus Complications;1(1999).
2. Alberti, KurtGeorgeMatthewMayer, and PZftZimmet. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and.
3. Alberti KG, Zimmet PZ. classification of diabetes mellitus. Provisional report of a WHO consultation. *DiabetMed*.1998;15(7):539-53. doi: 10.1002/(SICI)1096-9136(199807)15:7<539::AID-DIA668>3.0.CO;2-S, PMID 9686693.
4. Gress TW, Nieto FJ, Shahar E, Wofford MR, Brancati FL. Hypertension and antihypertensive therapy as risk factors for type 2 diabetes mellitus. Atherosclerosis Risk in Communities Study. *N Engl J Med*. 2000;342(13):905-12. doi: 10.1056/NEJM200003303421301, PMID 10738048.
5. AhmedAM. History of diabetes mellitus. *Saudi MedJ*.2002;23(4):373-8. PMID 11953758.
6. Zimmet PZ, McCartyDJ, de CourtenMP. The global epidemiology of non-insulin-dependent diabetes mellitus and the metabolic syndrome. & quot;J Diabetes Its Complications.1997;11(2):60-8. doi: 10.1016/s1056-8727(96)00090-6, PMID 9101389.
7. Association AD6.American Diabetes Association. Type 2 diabetes in children and adolescents. *Pediatrics*.2000;105(3):671-80. doi: 10.1542/peds.105.3.671.
8. Subaraman M, Gayathri R, PriyaVV. In vitro antidiabetic activity of crude acetone leaf extract of *Annonamuricata*. *Drug Invent Today*;14• Issue 6 • 2020.

9. Rajasekhar A, Peddanna K, Vedesree N, Munirajeswari P, Nagaraju N, Badri K Ret al. Antidiabetic activity of root tubers of *Asparagus gonoclados* Baker in streptozotocin induced diabetic rats. *J Ethnopharmacol.* 2019 Oct 5; 242: 112027. doi: 10.1016/j.jep.2019.112027. PMID 31226384.
10. Jangir RN, Jain GC. Evaluation of Antidiabetic Activity of hydroalcoholic extract of *Cassia fistula* Linn. pod in streptozotocin-Induced Diabetic Rats. *Pharmacogn J.* 2017; 9(5): 599-606. doi: 10.5530/pj.2017.5.95.
11. Dr. Radhika D, Priya R. Assessment of antidiabetic activity of some selected seaweeds. *ejbps.* 2015; 2(6): 151-4.
12. Abdul Vahab A, Harindran J. Antidiabetic activity of *Terminalia catappa* Bark extracts in alloxan induced diabetic rats; 16/05/2014.
13. Sikarwar MS, Patil MB. Antidiabetic activity of *Crateva nurvala* stem bark extracts in alloxan-induced diabetic rats. *J Pharm Bioallied Sci.* 2010 Jan-Mar; 2(1): 18-21. doi: 10.4103/0975-7406.62700, PMID 21814425.