



# International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)

IJAMSCR | Vol.12 | Issue 3 | Jul - Sept -2024

www.ijamscr.com

ISSN: 2347-6567

DOI : <https://doi.org/10.61096/ijamscr.v12.iss3.2024.366-379>



## Review

### Effectiveness of pictograms in patients or caregivers in healthcare settings: A systemic review

**P. Nikitha Reddy\***, V. Akash Goud, C. Mounika, B. Akhila, G. Neeraja, Syeda Madiha Faroohi

*Smt. Sarojini Ramulamma college of Pharmacy, Sheshadri nagar, Mahabubnagar-509001, Telangana, India*

\*Author for Correspondence: Dr. P. Nikitha Reddy  
Email: [patelnikithareddy6@gmail.com](mailto:patelnikithareddy6@gmail.com)

	<b>Abstract</b>
Published on: 2 Sept 2024	<b>Background:</b> Pictograms are a graphical symbol that convey's concept through its pictorial resemblance to a physical object, pictorial representation has shown to have a potential in enhancing patient knowledge.
Published by: DrSriram Publications	<b>Aim:</b> To study the effectiveness of pictorial health information on the patients or their caregivers.
2024  All rights reserved.  <a href="#">Creative Commons Attribution 4.0 International License.</a>	<b>Methodology:</b> The systematic review was conducted according to PRISMA guidelines. Randomized controlled trials including participants over 18 years older were included in the review. RCTs were included as it reduces certain type of bias by randomly allocating the participants. The risk of bias was assessed using Cochrane risk of bias tool, which assessed 5 different domains and scores were given to their indication of low, high and unclear risk. The interventions include in these articles were pictograms with text or pictograms with verbal instructions and low literacy plan pictograms. <b>Results:</b> A total of 965 articles were retrieved through electronic searching, which went through first and second pass screening. Among which articles were included for the system review. All the included studies show similar outcomes which said that pictograms had a positive impact on improving patient adherence to their respective medication and helped in decreasing dosing error. Factors such as age, gender, literacy level have a negative impact on adherence. Pictogram intervention improved adherence especially in patients with low health literacy levels than the written/oral interventions. <b>Conclusion:</b> The current review provided a brief literature on the effectiveness of pictogram in healthcare setting in patients or their caregivers of various age groups. Future studies should be aimed to identify the knowledge gaps and barriers impacting the effectiveness of pictogram for better patient education and safety. <b>Keywords:</b> Pictograms, Health literacy, Medication adherence

## INTRODUCTION

In developing countries, the consequences of disease conditions are rapidly increasing due to unhealthy lifestyles, physical inactivity, stress-full mind, and inadequate social and psychological well-being.<sup>1</sup> Drug resistance and unhygienic conditions, both communicable and non-communicable diseases are increasing extensively. These diseases cost an uncountable loss of lives a year and account for about 80% of difference in life expectancy worldwide.<sup>2</sup>

Prevention is a better way to keep diseases and remain healthy.<sup>3</sup>

Education is essential for the development of society; it not only helps in the development of the economy but plays a crucial role in the healthcare sector too. Inadequate understanding of healthcare leads to unsuccessful functioning in a pharmaceutical market designed for informed consumers.<sup>4</sup>

Health literacy is the ability to obtain, identify, determine, read, understand, and utilize the possibility of health-related information to make relevant decisions and follow up in medical treatments.<sup>5</sup> Individuals with adequate health literacy can take appropriate responsibilities on their health condition as well as their family's care.<sup>6</sup> Health related information can be provided in many ways which will help in development of health literacy among the population. The different methods such as icons or pictograms can be used of effectively improve the knowledge in these patients.<sup>7</sup>

A pictogram is a graphic symbol that conveys a concept through its pictorial resemblance to a physical object. Pictograms can surpass language as they can communicate speakers of many languages equally and effectively, even if language and culture are radically different.<sup>8</sup>

Pictograms consist of customized illustrations designed for giving health-related information, which includes indication, dosage form and their respective route of administration. It is not necessary that all the pictograms will be understood throughout every culture and different age groups, and among the people of low literacy level. In the matter of designing, lay participants are given a more active role called a "Pre-Designing Phase" which provides their inputs in the Designing of pictograms.

A pharmacist has a crucial role in medication history taking, drug education committees, therapeutic drug committees and integration of technologies.

Usually, medication leaflets and instructions for the use of medicines are written in high readability levels, which makes it difficult for the patients in the low health literacy population to adhere to the given medication. Pictogram is the best choice of tool for a better understanding of the drug-related information in patients with low health literacy [3]. Each drug information leaflet containing simple pictograms can be a useful tool in the enhancement of medication adherence and patient knowledge.

Even though the use of pictograms is potential in enhancing patient knowledge, there has been a lesser effort in evaluating the effect of these pictograms in the real-world population [10]. This study aimed to evaluate the effectiveness of the pictogram in improving patient knowledge and adherence to concomitant medication. Additionally, this systematic review will contribute to how positively the pictogram inclusion will affect the results of the intervention in low health literacy patients<sup>11</sup>

### Aim of this study

To study the effectiveness of pictorial health information on the patients or their care givers.

## MATERIALS AND METHODS

The following systematic review was carried out conferring to the preferred reporting items of the systematic reviews and meta-analysis [PRISMA] guidelines<sup>12</sup>.

The systematic review aimed at

1. Studying the effectiveness of health information on the patient.
2. To study the effectiveness of pictorial health information in low literacy people.
3. To study characteristics of pictorial health intervention used in healthcare
4. It helps us to understand if pictograms affected the increased medication adherence among the low literacy population.

### Inclusion and Exclusion criteria

Studies such as randomized controlled trial [RCT], blinded or open label were included for in the systematic review. RCTs following a cross-over design was excluded from the criteria. The study groups having more than one group were accommodated in the analysis.

The reason for inclusion was that RCTs helps to reduce a certain source of bias, accomplished by randomly allocated to two or more different groups, treated differently, and compared with a measured response. Articles other than RCTs including systematic reviews, observational studies, case reports, and narrative reviews were excluded.

Studies that included with pictogram or pictogram along with text fell under the inclusion criteria whereas studies without the inclusion of pictogram or on the phase of pictogram development were excluded.

Participants above 18 years old were included. The population included were of low health literacy.

### Search Methods for Identification of Studies


A systematic conduct of all available articles following randomized controlled trial design was Conducted. Studies aimed to find the effectiveness of pictograms on patient adherence and Knowledge was screened. Multiple keywords using appropriate Boolean operators were used to Build the search. The search was restricted to humans and the English language across the Databases.

Electronic Searches:

- I. PUBMED
- II. EMBASE
- III. CINAHL
- IV. SCOPUS

### Searching other Resources

We have hand searched many articles that have been included and relevant comments among the Information that can reclaim associated information. For advanced searches, the study database having disputes were identified in ongoing or unpublished trials.



### **Included Studies**

There were 15 RCT with pictograms as a comparator were included.

### **Study Design**

The 15 RCT (Zerfa 2014) consisted of parallel open label study {Browne 2018, Chan 2014, and Dowse 2005, Dowse 2014, Kher 2014, Mansoor 2006, Negradeh 2012, Phimarn 2018, Yin 2008}, double Blinded parallel group study {Kalichman 2013, Kriplani 2012}, single blinded parallel group study {Braich 2011, Murray 2007} and triple blinded parallel group study {Yin 2017}.

### **Interventions**

The intervention mostly included pictorial representation in different manner such as Simple PIL, text along with pictograms, low literacy plan with pictogram, verbal instruction along with Pictogram, standard and usual care along with pictograms.

### **Outcomes**

All the included study showed that similar outcome which stated that pictogram had a positive Outcome on improving the patient's knowledge adhere to their medication and helped in decreasing dosing errors. This study also distinguished other methods with pictogram and suggested that pictogram is indeed a better option for the healthcare management.

### **Excluded Studies**

965 articles were screened out of which 17 were excluded in the first screening, because it was found to be duplicate. In the second screening 151 articles were excluded based on title and Abstract. In the third screening 784 articles were excluded they were found to be irrelevant, wrong Intervention and negative outcome. In the end 15 articles, are mentioned based on title and abstract.

### **Data Extraction and Management**

The retrieved articles were screened simultaneously by two authors and in case of any disputes were solved by conversations with the other authors. The articles underwent a first-pass screening in which the title and abstracts were screened across the given inclusion criteria. Furthermore, the articles were analyzed through second pass screening where articles of full text was retrieved for inclusion of articles. Throughout the screening and data extraction process, two authors were involved simultaneously and encase of any accord was solved through consultation.

When multiple intervention groups were assembled, data retrieved from the different databases were combined to analyze for meaningful results. In the presence of multiple groups of participants, data from the group with the efficacy of pictogram better apprehended by the patient to that of a comparator were used in the review. Studies published in various parts, the primary article was used as a reference and secondary papers were necessary for deriving the additional information.

### **Assessment of Risk of Bias in Included Studies**

Using Cochrane risk of bias assessment tool two review authors independently assessed included articles.<sup>13</sup> The following domains were assessed and scored according to their indication of low (+), high (-) and unclear risk (?) were assigned.

- (i) Generation of allocation sequence;
- (ii) Allocation concealment;
- (iii) Blinding of participants, study personnel and assessors;
- (iv) Incomplete outcome data; and
- (v) Selective reporting.

Disputes in the studies were resolved by the discussion. The judgments behind each score reported in the table and assessment will be shown for individual study when combined in the figure.

### **Measures of Treatment Effect**

Number of events in the control and intervention groups of each study to calculate for odds ratio, percentile, variance, etc. were used in case of binary data whereas continuous data, the mean and standard deviations of individual study between the groups were identified. The authors ensured the consistency across the trials to avoid disputes. The 95% confidence interval (CI) was calculated for all outcomes. Considering the legitimate differences, the comparison of significance and guidance of the effect was reported by studies with their presentation.

### **Dealing with Missing Data**

When data for calculating odds ratio or mean were not accessible, the most advanced scientific data available that facilitated analyses of the included studies (e.g. test statistics, P values, etc.) was utilized. When such data was not possible (e.g. Measure of variation), values were imputed for the missing data by entering the comparable measure used from other pooled studies. Any discrepancies were exposed to sensitivity analysis.

### **Assessment of Heterogeneity**

To consider clinical and methodological characteristics of the included studies were adequately analogous for meta-analysis to contribute a clinically meaningful summary. The decision for pool studies were made by assessing the statistical heterogeneity by inspection of the Chi square test results and I<sup>2</sup> statistic. A rough guide to the interpretation of I<sup>2</sup> values is as follows:

- a) 0% to 40% might not be important;
- b) 30% to 60% may represent moderate heterogeneity;
- c) 50% to 90% may represent substantial heterogeneity; and
- d) 75% to 90% represents considerable heterogeneity.

These overlapping categories were considered, together with outcome uniqueness, in the assessment of heterogeneity.

### **Assessment of Reporting Bias**

In minimizing potential impact on reporting bias review authors did thorough examination of eligible studies or for any duplicated data present in database. The funnel plot for identifying the publication was not followed for the current review based on the decision by the authors.

### **Data Synthesis**

Most of the factors were likely to be influenced such as different hospital settings in different countries or difference in participant covariates. The data was combined using random-effects model together to form mixed effect model for population pharmacokinetics. Outcomes with continuous data were assessed for skew. When mean and SD were reported for studies, a rough check was made by determining observed mean minus lowest possible value and divided by Standard deviation. If the ratio was  $>1$ , it was considered that skew was likely. When skewed was considered likely outcome data were pooled, finding of each of these studies were included in the presentation of overall results for each outcome. (Appendix 1)

Study reference	country	Study design	N (loss to follow up)	Study population	intervention	control	Adherence effect measurement	result
Braich 2011 <sup>14</sup>	India	Multicentred, single-blinded, randomized control trail	225 (87)	Low literacy patients on cataract medication of eye drops of post operation	Medication use education: Pictograms in clinics; pictograms in home	Verbal instructions on medication use	Bottle amount measurement of eye drops at baseline and 28 days after surgery	Both intervention groups had significantly increased adherence as concluded by measuring percentage of eye drops used
Browne 2018 <sup>15</sup>	South Africa	Randomised control trial	116 (33)	Limited literacy HIV patients on ARVs	Medication information standard care with illustrated information	Standard care	Recognition by question at baseline, 1 month and 3-month post baseline interview	Significantly improved knowledge in intervention group of post baseline
Chan 2014 <sup>16</sup>	Malaysia	Randomised parallel, open-label study	126 (16)	Patients taking antihypertensive and anti diabetic medications	Medication labels; with enlarged fonts; and incorporating pictograms	Standard regular size to text medications labels without pictograms	MMAS-8 baseline and 4 weeks after intervention	No significant effect on the adherence between the study groups
Dowse 2005 <sup>[17]</sup>	South Africa	Randomised parallel, open-label study	87 (0)	Patients taking short course antibacterial medication	Medication labels; texts along with pictograms	Medication labels; Text only	Pill count/volumetric measurement and self-reporting of adherence after 3-5 days of intervention, combined adherence result	Significantly higher adherence with the intervention group as compared with the control group
Dowse 2014 <sup>[18]</sup>	South Africa	Randomised parallel, open-label study	116 (52)	HIV patients on new ARVT	PIL containing text along with pictogram	Standard care	Adherence self-efficacy scale of HIV treatment at baseline and 1,3, and 6 months after intervention	No significant difference on self-efficacy between the study groups

Kalichman 2013 <sup>[19]</sup>	USA	Randomised, double- blinded, parallel study	446 (45)	Low health literacy HIV patients on ARVT	Adherence counselling text along with pictograms, tool of choice of adherence: standard adherence of text along with illustrations comic strips and pill box of adherence tool	Counselling on general health improvement	HIV RNA load at baseline and 9 months after intervention and mostly PIL count from intervention start for 9 months	Patients with marginal health literacy had significantly greater undetectable HIV viral load in both intervention groups compared with the control and patients with low literacy didn't have significant effects on interventions
Kheir 2014 <sup>[20]</sup>	Qatar	Randomised, parallel, open label	123	Limited literacy skills in a culturally diverse multi-ethnic population	Verbal instructions along with text; and pictographic instruction only	Pictogram along with verbal instructions	Systematic approach of group discussions and interviews	Pictorials supported with verbal instructions was comprehended and pictogram only labels are least comprehending
Kripalani 2012 <sup>[21]</sup>	USA	Randomised, double blinded, parallel study	435	Patients with coronary heart disease	Post card reminders of refill; Illustrated schedule on medication with pictograms; combination of both	Usual care	Reported CMG of electronic pharmacy refill records for 1 year of follow up after intervention	No significant effect on adherence between the study groups
Mansoor 2006 <sup>[22]</sup>	USA	Randomised, parallel, open- label	127 (7)	Low health literacy patients on ARVT	PIL; with pictograms; and without pictograms	Usual care	PIL count and self- reported with questionnaires of approximate 14 days after intervention	Significantly increased adherence on PIL count and questionnaire of intervention compared with control group
Murray 2007 <sup>[23]</sup>	USA	Randomised single-blinded study	314 (44)	Heart failure patients $\geq 60$ years of age using cardiovascular medication	Pharmacist intervention: written information and containing pictograms	Usual care	Using prescription records and self- reported MEMS and MPR and questionnaires	Significant effect on overall adherence between intervention and control group in post intervention period. No significant effect on adherence between study groups of self-reporting

Negarandch 2013 <sup>[24]</sup>	Kurdistan	Randomised, open-label, parallel study	135 (8)	Low literacy patients with type 2 diabetes	Medication intervention education: teach back method: and with pictograms	Usual care	MMAS-8 at baseline 6 weeks post- intervention	Significantly higher adherence intervention group compared to control group. But no adherence between intervention groups
Phimarn 2018 <sup>[25]</sup>	Thailand	Randomised, parallel, open- label study	134	Low literate and poor adherence	Pictogram instructions	Traditional labels	Brainstorming, interviews and pilot evaluation	Experimental group had significantly higher post intervention understanding score than control group
Yin 2008 <sup>[26]</sup>	USA	Randomised, parallel, open- label study	245 (18)	Patients or caregivers of children on liquid medications	Medication counseling: with medication instruction sheets; pictograms: and teach-back	Usual care	Self-reported adherence by interview at 3-5 days after medicine dispensing	Significantly higher adherence in the intervention group with the control group
Yin 2017 <sup>[27]</sup>	USA	Randomised, parallel, Triple-blinded study	259 (42)	Low literate patients of children with asthma	Asthama action plan: of low literacy plan along with pictogram	Standard care plan	Questionnaire to assess error in knowledge	Intervention group had significantly higher impact than control group
Zerafa 2011 <sup>[28]</sup>	Malta	Randomised control trial	86 (6)	Cardiac surgery patients on medication	Pharmacist intervention: with written medication information sheets and pictograms	Usual care	Questionnaire of patient compliance after 8 weeks after surgery discharge	Significantly higher patient compliance in intervention group with control group



## RESULTS

### Search

Electronic search was conducted in PubMed, EMBASE, CINAHL and SCOPUS that yielded a total number of hits 241, 98, 113 and 515 respectively. A total of 965 articles were identified through this search. Out of 965 articles identified, 17 articles were excluded since they were found to be duplicates. In the second screening, 151 articles were excluded out of 948 articles based on the title and abstract screening. 782 articles were further excluded as outcome, irrelevant and intervention as exclusion criteria. Finally, 15 articles were included in the systematic review.

### Risk of Bias

All the studies included<sup>15</sup> reported adherence to pictogram-based interventions (Appendix 2). The patient groups were mostly who were on antiretroviral medications, cardiac patients, and patients on post-operative cataract medication, patients and care givers with less health knowledge. Quality of the studies were evaluated by two reviewers using Cochrane collaboration Tool.

Significant studies were identified as open-label studies (93.3%, High risk). In two studies blinding of participants and key study personnel were ensured (13.3%, low risk). Thirteen trials (86.6%) were reported as high risk of detection bias. Five studies reported an unclear risk of selection bias (33.3%, Allocation concealment). One study reported high risk of attrition bias and 3 studies (20%) reported unclear risk. No study reported high risk or unclear risk of selection bias (random generation). As seen in figure 2, all the studies had at least one dimension with a high risk of bias, but met the acceptable quality.

In this study, we assessed various articles to evaluate the pictograms in patient understanding and medication adherence. From the 15 included studies 3 studies done by Chan<sup>16</sup>, Dowse<sup>18</sup>, Kripalani<sup>21</sup> did not have a statistically significant pictogram effect on medication adherence. Each study varied with the use of pictograms. In study interventions, pictograms were used alone or in combination with the text-based/written or verbal/oral instructions of medication use. The current review points of the conclusion of improved patient adherence were with the combinational use of pictograms with text-based and/or verbal instructions.

A few article reviews resulted in an insignificant effect that were attributed to insufficient data, sample size or required further studies to have a significant conclusion. The adherence was assessed by questionnaire, self-reporting and interviews. Significant evidence was there to conclude that pictogram-based interventions would enhance the medication adherence of patients.

The interventional complexity acts as a limitation for the pictogram contrition to medication adherence. Measuring adherence was difficult to evaluate and required standardization methods.


In this review, a study by Kalichman et al<sup>19</sup>, used pictogram intervention along with adherence counselling of text with a pictogram, and adherence tools of choice of illustration with comic strips and pillbox. So, it was difficult to conclude the adherence effect of pictogram alone from the adherence counselling. The RCT conducted by Negrandeh et al<sup>24</sup>, was done in a diabetic clinic by a nurse among low health literacy patients with type 2 diabetes. The educational medication intervention was through teach back method and pictograms against the usual care as the control group. This resulted with no adherence between the intervention groups and significantly higher adherence in the intervention group as compared with the control group.

Whereas, a study led by Mansoor et al<sup>22</sup>, was done on low health literacy patients on antiretroviral therapy (ART) with patient information leaflet (PIL) and without pictograms against the standard care. A significant increase in adherence to patients receiving PIL with pictograms as compared with other groups. Study designs of comparison of interventional groups show only the difference in the use of pictograms. Patient-related factors play a key role in the contribution of medication adherence. Factors such as age, gender, literacy level have a negative impact on adherence. Pictogram intervention improved adherence especially in patients with low health literacy levels than the written/oral interventions. RCT study by Chan<sup>16</sup>, Dowse<sup>18</sup>, Kripalani<sup>21</sup>, Murray<sup>23</sup>, Yin<sup>27</sup> and Zerafa<sup>28</sup> doesn't discuss about the role of health literacy for medication adherence. It indicated that the pharmaceutical pictogram is most beneficial for the patients challenged with low health literacy.

Another known factor influencing medication adherence was the nature of the therapy. The RCT study by Browne<sup>15</sup>, Mansoor<sup>22</sup>, Kalichman<sup>19</sup>, and Dowse<sup>18</sup> was on ARVT; Chan<sup>16</sup> was on antihypertensive and anti-diabetic medications; Murray<sup>23</sup> was on cardiovascular medications.

But the above-mentioned articles did not specifically describe about prescribed therapies of the included participants. Therapeutic effects are based on dosing and frequency of therapy, patient's attitude, belief, adverse events of the treatment, and effectiveness of the medication. However, the studies performed by Browne<sup>15</sup>, Mansoor<sup>22</sup>, and Murray<sup>23</sup> stated a significant effect on pictogram-based interventions. The adherence effect was measured by different methods in various studies. Such included the bottle amount measurement, questionnaire self-report interview, pill count, and group discussion, electronic refill records by pharmacy, brainstorming. There was no single method recommended, and so the studies use a mixed method for measuring the adherence.

The RCT study by Yin et al<sup>26</sup>, review the pictogram effects on caregivers in the administration of liquid medication and suggested that it may result in the reduction of dosing error with enhanced comprehension and improved adherence.



	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)
Braich 2010	+	?	-	-	+	+
Browne 2018	+	+	-	-	-	+
Chan 2014	+	?	-	-	+	+
DROWSE 2004	+	?	-	-	+	+
Drowse 2014	+	+	-	-	?	+
Kalichman 2013	+	+	+	-	?	+
Kheir 2014	+	?	-	-	+	+
Kripalani 2012	+	?	-	-	?	+
Mansoor 2006	+	+	-	-	-	+
Murray 2007	+	+	-	-	+	+
Negrandeh 2013	+	+	-	-	?	+
Phimarn 2018	+	?	-	-	+	+
Yin 2008	+	+	-	-	+	+
Yin 2017	+	+	-	-	+	+
ZERAFA 2011	+	?	-	-	+	+

## DISCUSSION

To understand information about one's own medication is vital for each patient for better medicine use and safety. Various literacy rates across the globe are a potential barrier in the interpretation of written information. The health literacy of patients provides relevance in the development of information tools to provide adequate understanding for such populations<sup>14</sup>. Pharmaceutical

Pictograms are a remarkable substitute unless they are not sensitive socially. A various study has tested the effectiveness of pictogram in various settings to identify its usefulness in improving patients' understanding amongst various literacy groups<sup>29</sup>. Pictograms play an evident role in transforming the medical information to improve understanding, adherence and in medication recalling. Even though various forms of pictograms have been developed and tested, the efficacy remains questionable in improving medication related parameters. This can be overcome by providing dedicated patient counselling utilizing the pictograms for better medication use. A validated model for the design and interpretation of pictograms are the need for the hour for imparting health information and for providing better patient safety<sup>30</sup>. Pictorial aids are also effective in caregivers to assist in the administration of certain dosage forms of medications.

Pictograms use will improve the understanding of medication instructions, the dosing accuracy, and also will improve recall information in the caregivers for better patient care. Incorporating pictograms into verbal instructions or counselling on medications or the text instructions was more beneficial than to be used alone. The health literacy level of caregivers will also get contributed by pictorial aid effectiveness<sup>31</sup>. Pictograms will recall, enhance comprehension and adherence with medication information.

The pictograms have the ability in helping the population with low literacy, remains to be a concern, considering the low education level and socioeconomic status of the patient population. Successful establishment of medication safety programs are essential for the development of healthcare setting through the reduction of cost and for delivering better patient care. The increase in rate of right interpretation of pictograms can be of utmost use whenever provided as a replacement to instructions in verbal form. The impact of pictograms in preventing medication administration errors in a health care setting and in improving clinical outcomes needs to be reviewed and studied spontaneously to explore future outcomes of pictograms.

The current systematic review aimed to investigate about the potential effectiveness on patient of medication adherence towards the therapies based on collecting and summarizing the shreds of evidence depending on certain inclusion and exclusion criteria of pictograms. It is usually hard for patients to retain verbally communicated information, for which this short systematic review conducted by Van Beusekom 2018 to evaluate the extent and effects of patient involvement in the design and development of the pictograms for written information of drug. This review included 73 articles which were published between 1993 and 2018, this review focused on two groups, one being the patient party and the other being the non -patient party that helped in the design process of the pharmaceutical pictograms. It showed that the involvement in the design process of lay participants led to pictograms that were preferred in the specific target group. The involvement of lay participants also showed to consistently lead a positive effect on the terms of patient understanding. Overall this review showed involvement of the non-participants in the development of pharmaceutical pictograms and provided evidences involving lay end-users in the design process to help to increase the likelihood that resulting pictograms are well-understood, well-received and aid recalls of the drug information that supports. It is also essential to involve participants that meet the key criteria in the targeted group in the evaluation of pictograms and pictogram based information as it was seen from that different audiences perceive information differently.<sup>32</sup> Similarly, the current review was based on Pictogram alone and pictogram with text as the major intervention. The results of the current study gave a mixed perception of the patients and their care givers in understanding the effectiveness of pictogram.

Another review led by Sletvold 2019 and team focused on the impact of pictograms on medication adherence. This study initially included 1283 articles, out of which only 17 articles were included for analysis after excluding the others based on duplication and the inclusion criteria. The study population was diverse in clinical disorders, treatment regimen, terms of age and the level of health literacy. of the included studies 10 articles reported a statistically significant effect of pictograms used in the studies varied though most of them used pictograms along with text based or verbal instructions. Patient-related factors such as age, long term medication use, and a different type of disease population also affect medication adherence which is why it is necessary to develop pictograms that are focused on the target population. It was concluded that pictograms do serve as a communication tool in combination with verbal or oral instruction to enhance visual attention, comprehension, recall, and adherence to medications.<sup>33</sup> In context to the above study, our study search was conducted till 2019 September and few articles were added which increased the sample population of the present review to give more vivid results. A systematic review based on Magnay 2018 was concerned with the validation process or development of methods for assessing menstrual blood loss which was based on the different strategies used to develop NICE guidelines.

1438 records were retrieved out of which 71 fulfilled the inclusion criteria, which was used to determine methods to measure the Menstrual Blood Loss (MBL) and to distinguish between the normal and Heavy Blood

Flow (HBL), the suitable diagnosing for HMB and routine clinical practice and practical and limitations to research background settings. This review showed that every available method cannot assess MBL. A pictorial representation showed a balance in ease of understanding and validated pictograms in MBL determination in each way using QOL in both clinical and research settings.<sup>34</sup> This study focused on a single population and condition whereas the present study had various intervention and the target population varied among the studies. The interventions were mainly focused on low literacy population and the intervention were keenly designed to improve the adherence and the knowledge in the specific population.

Another review conducted by Chan, 2015 included studies that used pictorial aids with liquid medication and measured its dosing accuracy, comprehension of instruction, recall information and adherence of caregivers. 1363 records were yielded from the search out of which only 5 studies met the search which contained 962 participants, a wide range of liquid formulations were studied including prescription and OTC drugs. Regarding dosing errors, pictograms were given to one half of the population and the other half received text information and it was seen that the group which received pictograms showed fewer mistakes in dosing error. As for the recall of medication, it was also done the same way, one half received pictograms while the other half received the non-pictogram intervention. The group that received pictograms recalled their medication instructions better as compared to the other group. When all the criteria were combined it was seen that pictorial aids are useful intervention based on the findings. The study had a direct comparison analysis.<sup>35</sup> The study intervention included only pictogram as intervention whereas our study was based on the pictogram alone or with some amount of text with the same. The outcomes measures were medication adherence more precisely than other outcomes relating to dosing and dosage forms.

Only a few studies among the included studies were reviewed for dosage forms where pictograms were efficiently used and served the purpose.

### Limitations

A systematic review included 15 studies. Each of the studies were randomized controlled trials. The trials differed based on the design of blinding where few studies were blinded, and many were open label. This led to high amount of heterogeneity across the studies. The outcomes were reported in various data forms which was a drawback in conducting a meta-analysis. The intervention yielded a mixed response since the pictogram and text were utilized. Future studies focusing on pictograms alone and outcome measured using a uniform tool should be targeted for the design of meta-analysis to determine the effectiveness of pictogram in various healthcare settings

## CONCLUSION

Pictograms have been an essential tool for educating patients of various literacy groups. The effectiveness of pictograms remains unclear regardless of the group studied. The current review provided a brief literature on the effectiveness of pictogram in healthcare setting in patients or their caregivers of various age groups. Future studies should be aimed to identify the knowledge gaps and barriers impacting the effectiveness of pictogram for better patient education and safe.

## REFERENCES

1. Murray CJ, Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. *The Lancet*. 1997; 349(9061):1269-76
2. Garrett L. Existential challenges to global health. New York: Center on International Cooperation, New York University. 2013.
3. Klein RE, Weller SC, Zeissing R, Richards FO, Ruebush TK. Knowledge, belief and practices in relation to malaria transmission and vector control in Guatemala. *Am J Trop Med. Hyg*. 1995; 52(5):383-8.
4. Parker RM, Ratzan SC, Lurie N. Health literacy: a policy challenge for advancing high-quality health care. *Health Aff*. 2003; 22(4):147-53.
5. Hewitt M. Roundtable on Health Literacy; Board on Population Health and Public Health Practice; Institute of Medicine. Facilitating State Health Exchange Communication Through the Use of Health Literate Practices: Workshop Summary. National Academies Press. p. 1; 2012
6. McQueen DV, Kickbusch I, Potvin L, Pelikan JM, Balbo L, Abel T. Health and modernity: the role of theory in health promotion. Springer Science & Business Media; 2007.
7. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Viera A, Crotty K. et al., Health Literacy Interventions and Outcomes: An Updated Systematic Review. *Evid Rep Technol Assess*. 2011; 199:1-941.
8. Health Literacy and Patient Education Guide: Pictograms. Strauss Health Science Library. Last update 2020.
9. Schroyens W, Schaeken W, Handley S. In search of counter-examples: Deductive rationality in human reasoning. *Q J Exp Psychol A*. 2003 1; 56(7):112945.

10. van Beuskom M, Kerkhoven AH, Bos MJW, Guchelaar HJ, van den Broek JM. The extent and effects of patient involvement in pictogram design for written drug information: a short systematic review. *Drug Discov Today*. 2018; 23(6):1312-1318.
11. Chan HK, Hassali MA, Lim CJ, Saleem F, Tan WL. Using Pictogram to assist caregivers in liquid medication administration: Systematic review. *J Clin Pharm Ther*. 2015; 40(3):266-72.
12. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009; 151(4):264-9, W264.
13. Higgins JPT, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD et al. The Cochrane Collaboration's tool for assessing risk of bias in Randomised trials. *BMJ*. 2011; 343:d5928.
14. Braich PS, Almeida DR, Hollands S, Coleman MT. Effects of pictograms in educating 3 distinct low-literacy populations on the use of postoperative cataract medication, *Can J Ophthalmol*. 2011; 46(3):276-81.
15. Browne SH, Barford K, Ramela T, Dowse R. The impact of illustrated side effect information on understanding and sustained retention of antiretroviral side effect knowledge. *Res Social Adm Pharm*. 2019; 15(4):469-73.
16. Chan HK, Hassali MA. Modified labels for long-term medications: influences on adherence, comprehension and preferences in Malaysia. *Int J Clin Pharm*. 2014; 36(5):904-13.
17. Dowse R, Ehlers M. Medicine labels incorporating pictograms: do they influence understanding and adherence? *Patient Educ Couns*. 2005; 58(1):63-70.
18. Dowse R, Barford K, Browne SH. Simple, illustrated medicines information improves ARV knowledge and patient self-efficacy in limited literacy South African HIV patients. *AIDS Care*. 2014; 26(11):1400-06.
19. Kalichman SC, Cherry C, Kalichman MO, Amaral C, White D, Grebler T, et al. Randomized clinical trial of HIV treatment adherence counselling interventions for people living with HIV and limited health literacy. *J Acquir Immune Defic Syndr*. 2013; 63(1):42-50.
20. Kheir N, Awaisu A, Radoui A, El Badawi A, Jean L, Dowse R. Development and evaluation of pictograms on medication labels for patients with limited literacy skills in a culturally diverse multiethnic population. *Res Social Adm Pharm*. 2014; 10(5):720-30.
21. Kripalani S, Schmotzer B, Jacobson TA. Improving medication adherence through graphically enhanced interventions in coronary heart disease (IMAGE-CHD): a randomized controlled trial. *J Gen Intern Med*. 2012; 27(12):1609-17.
22. Mansoor LE, Dowse R. Medicines information and adherence in HIV/AIDS patients. *J Clin Pharm Ther*. 2006; 31(1):7-15.
23. Murray MD, Young J, Hoke S, Tu W, Weiner M, Morrow D, et al., Pharmacist intervention to improve medication adherence in heart failure: a randomized trial. *Ann Intern Med*. 2007; 146(10):714-25.
24. Negarandeh R, Mahmoodi H, Noktehdan H, Heshmat R, Shakibazadeh E. Teach back and pictorial image educational strategies on knowledge about diabetes and medication/dietary adherence among low health literate patients with type 2 diabetes, *Prim Care Diabetes*. 2013; 7(2):111-18.
25. Phimarn W, Ritthiya L, Rungsoongnoen R, Pattaradulpithuk W, Saramunee K. Development and Evaluation of a Pictogram for Thai Patients with Low Literate Skills. *Indian J Pharm Sci* 2019; 81(1):89-98
26. Yin HS, Dreyer BP, van Schaick L, Foltin GL, Dinglas C, and Mendelsohn AL. Randomized controlled trial of a pictogram-based intervention to reduce liquid medication dosing errors and improve adherence among caregivers of young children. *Arch Pediatr Adolesc Med*. 2008; 162(9):814-22.
27. Shonna Yin H. Health Literacy and Child Health Outcomes: Parental Health Literacy and Medication Errors. *Health Literacy and Child Health Outcomes*. 1938
28. Zerafa N, Adami MA, Galea J. Impact of drugs counselling by an undergraduate pharmacist on cardiac surgical patient's compliance to medicines. *Pharm Pract*. 2011; 9(3):156-61.
29. Montagne M. Pharmaceutical pictograms: a model for development and testing for comprehension and utility. *Res Social Adm Pharm*. 2013, 9(5), 609-20.
30. Knapp P, Raynor DK, Jebar AH, Price SJ. Interpretation of medication pictograms by adults in the UK. *Ann Pharmacother*. 2005; 39(7-8):1227-33.
31. Morrell RW, Park DC, Poon LW. Effects of labelling techniques on memory and comprehension of prescription information in young and old adults. *J Gerontol*. 1990, 45(4): 166-72.
32. van Beusekom MM, Kerkhoven AH, Bos MJW, Henk-Jan Guchelaar, van den Broek JM. The extent and effects of patient involvement in pictogram design for written drug information: a short systematic review. *Drug Discov Today*. 2018; 23(6):1312-18.
33. Sletvold H, Bjørnli Sagmo LA, Torheim EA. Impact of pictograms of medication adherence: A systematic literature review. *Patient Educ Couns*. 2019, S0738-3991(19)30562-2.

34. Magnay JL, O'Brien S, Gerlinger C, Seitz C. A systematic review of methods to measure menstrual blood loss. *BMC Women's Health*, 2018; 18:142.
35. Chan HK, Hassali MA, Lim CJ, Saleem F, Tan WL. Using pictograms to assist caregivers in liquid medication administration: a systematic review. *J Clin Pharm Ther*, 2015; 40(3):266-72