



International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)

IJAMSCR | Volume 3 | Issue 2 | April-June- 2015
www.ijamscr.com

ISSN: 2347-6567

Research article

Medical research

Awareness of antenatal exercises among pregnant women in tertiary care centers, Mangalore, India

Rachana Nayak, Simeen Virani, Chetna Gupta, Vijaya kumar K*, Dr. Amitesh Narayan #
Dr. Suchitra Thunga \$, Dr. Prasanna Mithra P. §

*Associate Professor, Department of Physiotherapy, Kasturba Medical College, Mangalore, Manipal University, Karnataka, India.

#Professor and Head, Department of Physiotherapy, Kasturba Medical College, Mangalore, Manipal University, Karnataka, India.

\$Professor, Department of Obstetrics and Gynaecology, Kasturba Medical College, Mangalore, Manipal University, Karnataka, India.

§Associate Professor, Department of Community Medicine, Kasturba Medical College, Mangalore, Manipal University, Karnataka, India.

*Corresponding author: Vijaya kumar K

Email: vijay.kk@manipal.edu

ABSTRACT

A proper exercise regimen during antenatal care helps the mother have a safe pregnancy and delivery. A physiotherapist trained in maternal health is best suited to prescribe exercises to pregnant women. In this cross sectional study, the participants were recruited from KMC hospitals, Mangalore from December 2014-February 2015. Pregnant women in their reproduction age between 18 – 40 years in any trimester who had visited for antenatal check-up in Department of Obstetrics and Gynaecology were included in our study with the aim to find out their level of awareness about Antenatal Exercises in Pregnancy. A self-administered questionnaire was developed from previous literatures which was reviewed and content validated by subject experts in their respective fields. A sample of Ninety pregnant women who met the study criteria were explained about the study. After obtaining their Informed Consent the antenatal awareness questionnaire were given to pregnant women either Kannada or English depending on their language preference. The questionnaire includes basic demographic data, pregnancy history and question to relate to antenatal classes, different types of exercises prescribed during pregnancy. Data was entered and analyzed using SPSS version 17. Frequencies and percentages were taken out as part of descriptive statistics

Keywords: Antenatal exercises, Pregnancy, Obstetrics and Gynecology.

INTRODUCTION

Physical activity is a very important aspect of good health. Exercise is very important irrespective of age and especially important for pregnant women as it

helps them overcome pregnancy related complications and maintain good physical fitness. Exercise during pregnancy not only helps in maintaining musculoskeletal fitness but also helps in controlling weight, maintaining blood glucose, to

cope with various psychological and physical stresses during pregnancy and labour, strengthening muscles for labour and improves blood circulation.^[1,2] Kegel's exercise is a commonly followed prenatal exercise used to strengthen pelvic floor muscle which may stretch during delivery, hence prepare for labour.^[3,4,5] To cope with emotional stresses and labour pain, breathing techniques and relaxation techniques are also proved to be successful.^[6] Ante natal exercise also decreases adipose tissue growth, increases stress tolerance and advances neuro behavioural relaxation in the foetus.^[7] Various exercise regimen can be followed to improve overall fitness. ACOG recommends low impact moderate intensity exercise for pregnant women, which can be gradually progressed over a period of time.^[8] One of the commonly recommended programme includes aerobic exercise which affects heart rate, circulatory system. A proper exercise regimen during antenatal care helps the mother have a safe pregnancy and delivery.^[9] A physiotherapist trained in maternal health is best suited to prescribe exercises to pregnant women. There have been studies in various developed and developing countries to assess awareness of benefit of exercise during pregnancy.^[10-13] However there are limited literature to know about awareness of antenatal exercises among pregnant women from India.

METHODOLOGY

The study protocol was approved by Institutional Ethics Committee of Kasturba Medical College, Mangalore, Manipal University, India. In this cross sectional study, the participants were recruited from KMC hospitals, Mangalore from December 2014-February 2015. Pregnant women in their reproduction age between 18 – 40 years in any trimester who had visited for antenatal check-up in Department of

Obstetrics and Gynecology were included in our study with the aim to find out their level of awareness about Antenatal Exercises in Pregnancy. A self-administered questionnaire was developed from previous literatures^[11,12] which was reviewed and content validated by subject experts in their respective fields. A sample of Ninety pregnant women who met the study criteria were explained about the study. After obtaining their Informed Consent the antenatal awareness questionnaire were given to pregnant women either Kannada or English depending on their language preference. The questionnaire includes basic demographic data, pregnancy history and question to relate to antenatal classes, different types of exercises prescribed during pregnancy. Data was entered and analyzed using SPSS version 17. Frequencies and percentages were taken out as part of descriptive statistics

RESULTS

The demographic variables in our study are 83% of the respondents were above 25 years of age, 46% had completed graduation/ post-graduation in their education level. Their socio economic status was assessed on the Kuppuswamy scale and it was found that 34% of our respondents are under the upper class and 66% are under upper middle class. 72% samples in our study were multiparous. The results are presented in Table 1. 46% of study participants were aware of antenatal classes and the main source of awareness from family/friends are reported by 48% and followed by 30% attended the antenatal class. Only 20% of them aware of different type of exercises in antenatal care. The results are presented in Table 2. Awareness of Antenatal Exercises among Pregnant Women in Tertiary Care Centers, Mangalore

Table 1 Demographic Characteristic of the Respondents

Variables	N (%)
Age	
<25 years	15 (17)
25 and above	75 (83)
Education level	
High School Certificate	21 (23)
Intermediate / Post high school diploma	28 (31)
Graduate / Post graduate	41 (46)

Occupation	
Unemployed	23 (26)
Clinical / Shop owner / Farmer	27 (29)
Semi profession	08 (09)
Profession	32 (36)
Socio economic status	
Upper	31 (34)
Upper Middle	59 (66)
Parity	
Primi parous	25 (28)
Multiparous	65 (72)

Table 2 Awareness of Antenatal Classes and Exercises among Respondents

Variables		N (%)
Awareness of antenatal classes	Yes	41 (46)
	No	49 (54)
If yes how	Family /Friends	20 (48)
	Media	07 (17)
	Health care centers	02 (05)
	Attended the Antenatal class	12 (30)
Are you Aware Of these Exercise in Antenatal care Aerobics	Yes	8 (09)
	No	63 (70)
	Not Sure	19 (21)
Back Care Exercises	Yes	18 (20)
	No	52 (58)
	Not Sure	20 (23)
Abdominal Exercises	Yes	19 (21)
	No	52 (58)
	Not Sure	19 (22)
Pelvic Muscle Exercises	Yes	12 (13)
	No	58 (65)
	Not Sure	20 (22)
Relaxation with breathing Exercises	Yes	19 (21)
	No	55

		(61)
	Not Sure	16
		(18)
Muscle strengthening/ Balance Exercises	Yes	17
		(19)
	No	55
		(61)
	Not Sure	18
		(20)
Are you Referred to Physiotherapy for antenatal exercises during present pregnancy	Yes	12
		(13)
	No	78
		(87)
	Back care exercise	06
If yes, what Exercises were prescribed	Pelvic exercise	03
	Relaxation	03

DISCUSSION

Antenatal care is an essential pillar of safe motherhood. To the best of our knowledge, this study is the first of this kind to know the level of awareness of antenatal exercises among pregnant women from India. In our study, only 46% of respondents are aware of Antenatal classes. Among the main source of information about antenatal exercises, 48% was from Family /Friends, whereas 30% attended the Antenatal Class and 17% from Social media. A recent study from Sarfaraz et al from Pakistan reported that 83.7% of pregnant women were aware of antenatal exercises, 42% attended the antenatal class, and the main source of information was from health care centers (48.8%). Though in our study, the 50% participants were graduates and also upper middle class, only 5% of the participants reported that health care centers were the source of information for Antenatal Exercise. When the respondents were asked about the different type of exercises in antenatal care, only 9% were aware of the aerobic exercises, 21% for back care, 21% for abdominal exercises, 13% for pelvic floor exercises, breathing and relaxation techniques, and 19% for muscle strengthening and balance training exercise, which states that our pregnant women were the poor knowledge about the antenatal exercises during pregnancy. A recent study from Mbada EC et al had reported that where 59% were aware about aerobic exercises, 70% for pelvic floor exercise, 97% for muscle strengthening exercises, 56% for abdominal exercises, and 97% for back care exercises. Our study finding shows that participants were poor awareness

about the role of different exercises and benefits for antenatal care. Physical therapy plays a major role in antenatal care, however, in our study, only 13% of respondents were referred for physiotherapy during antenatal care, mainly for back care exercises. Performing all or combination of different exercises under the supervision of a Physical therapist results in various benefits. Hence, there is a need to incorporate Physiotherapy during routine antenatal care.

LIMITATIONS

The outcome of this study is limited in its generalizability and needs to be validated in other settings. The study is also limited to two months period. Our community is a multilingual community (Malayalam, Tulu, Konkani), however, we limited our samples to those who know either English or Kannada and also upper socioeconomic class of the society.

CONCLUSION

The important findings that emerge from this study in Mangalore are that awareness about antenatal exercises is inadequate among pregnant women and the role of physiotherapy interventions in antenatal care among health professionals. These can be achieved through awareness campaigns, physiotherapy health education, and interaction with the health professional.

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interests regarding the publication of this paper.

ACKNOWLEDGMENTS

The authors gratefully acknowledge all the women who volunteered for this study. Also, they are

grateful to the administrative and nursing staff of the hospitals.

REFERENCES

- [1]. Wolfe LA, Davies GAL. Canadian guidelines for exercise in pregnancy. *Clin Obstet Gynecol.* 2003; 46:488–95.
- [2]. Morris SN, Johnson NR. Exercise during pregnancy: a critical appraisal of the literature. *J Reprod Med.* 2005; 50:181–8.
- [3]. Park, S.H., Kang, C.B., Jang, S.Y. and Kim, B.Y. (2013) Effect of Kegel Exercise to Prevent Urinary and Fecal Incontinence in Antenatal and Postnatal Women: Systematic Review. *Journal of Korean Academy of Nursing*, 43, 420- 430.
- [4]. El Hamid, A.A., Azzam, H.F., Ismail, G.M. and Gaafar, H.M. (2012) Effect of a Structured Antenatal Kegel Exercises Protocol on Labor Progress among Women Attending Antenatal Clinics. *Egyptian Nursing Journal*, 3, 2090-210.
- [5]. Dumoulin C, Hay-Smith J. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Cochrane Database Syst Rev.* 2010 Jan (1):CD005654.
- [6]. Green JM. Expectations and experiences of pain in labor: findings from a large prospective study. *Birth* 1993; 20: 65- 72.
- [7]. Clapp JF. Exercise during pregnancy. A clinical update. *Clin Sports Med.* 2000; 19:273–86.
- [8]. ACOG Committee Obstetric Practice. ACOG Committee Opinion. Number 267, January 2002: exercise during pregnancy and the postpartum period. *Obstet Gynecol.* 2002; 99:171– 173.
- [9]. Kramer MS, McDonald SW. Aerobic exercise for women during pregnancy. *Cochrane Database Syst Rev.* 2006 Jul 19; (3):CD000180.
- [10]. Ribeiro CP, Milanez H. Knowledge, attitude and practice of women in Campinas, São Paulo, Brazil with respect to physical exercise in pregnancy: a descriptive study. *Reprod Health.* 2011; 8(1):31.
- [11]. Sarfraz M, Islami D, Hameed U, Hasan Danish S, Ahmad F. Role of Physical Therapy in antenatal care as perceived by the clients-a cross sectional survey on pregnant females attending antenatal OPD. *Pakistan Journal of Medicine and Dentistry* 2013; 1(01): 34-46.
- [12]. Olufemi M, Florence A, Bengami O. Perception of pregnant mothers attending antenatal clinic on usefulness of prenatal exercise in Osogbo, Orun State Migeua. *Health Science Journal.* 2014: 8(2).
- [13]. Mbada EC, Adebayo EO, Adeyemo BA, Arije O, Dada O. Knowledge and Attitude of Nigeran Pregnant women towards ante natal exercises: A cross sectional survey. *ISRN Obstetrics and Gynecology.* 2014; 1-8.