International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)
IJAMSCR |Vol.12 | Issue 3| Jul - Sep -2024
www.ijamscr.com
ISSN: 2347-6567
DOI : https://doi.org/10.61096/ijamscr.v12.iss3.2024.311-319

Review

COLLAFIBE SACHETS: An Advanced Liposomal Technology Encapsulated with pepzyme pro to boost collagen peptide Absorption to support hydration & Elasticity of skin.

GovindShukla[1], Dr. Chandramauli[2], Dr. Balaswamy N.G[3], Dr. Rajkumar[4], C.J. Sampath Kumar [5]

Pugos Nutrition Research Centre Hyderabad , A unit of PUGOS Products Pvt. Ltd. 42, 2nd Floor, Leelavathi Mansion, 6th Cross, Margosa Main Road Malleshwaram Bangalore-56003, India.

*Author for Correspondence: GovindShukla
Email: lactonovaresearch44@gmail.com

Published on: 22 Jul 2024

Published by:
DrSriram Publications

2024| All rights reserved.

Creative Commons Attribution 4.0 International License.

Abstract

It is evident that marine collagen in collafibe sachets is a versatile compound capable of healing skin injuries of varying severity, as well as delaying the natural human aging process. From in vitro to in vivo experiments, collagen has demonstrated its ability to invoke keratinocyte and fibroblast migration as well as vascularization of the skin. Additionally, marine collagen and derivatives have proven beneficial and useful for both osteoporosis and osteoarthritis prevention and treatment. Other bone-related diseases may also be targeted by collagen, as it is capable of increasing bone mineral density, mineral deposition, and importantly, osteoblast maturation and proliferation. Marine organisms harbor numerous bioactive substances .Scientific research on various applications of collagen extracted from these organisms has become increasingly prevalent. Marine collagen in collafibe sachets can be used as a biomaterial because it is water soluble, metabolically compatible, and highly accessible. The present Article Reviews the role of collafibe sachets in Improving cell turnover and collagen formation, which keeps your skin stay elastic and healthy.

Keywords: Collafibe Sachets, Glowing Skin, Healthy Hairs, Nails

INTRODUCTION

The UVR that affects the skin is composed of two types of waves: UVA and UVB. UVB rays are shorter than UVA rays, and are the main cause behind inflammation and melanin production. However, it is the UVA rays, with their longer wavelength, that are responsible for much of the damage associated with photoaging. UVA rays penetrate deep into the dermis, where they damage collagen fibers, leading to wrinkle formation.



Technical advancements in collafibe sachets
Liposomal Technology: Vitamin C and Glutathione
Pepzyme Pro boosted Marine Collagen with Peach Extract,
Cold Water Soluble Vitamins



Fig 1: Reduces wrinkles & increase elasticity

UV rays induce the production of in situ radical oxygen species (ROS) and matrix metalloproteinases (MMP). These factors are the root of wrinkle formation because they destroy the collagen matrix in the dermis. Fortunately, the skin's repair mechanism will rebuild the damage collagen. However, the hindrance of skin renewal by repeated exposure to uncontrolled levels of ROS and MMP leads to the formation of wrinkles. The extracellular matrix (ECM) plays important roles in the physical integrity of cells, where it is involved in cell proliferation, differentiation, migration, and adhesion [1–6]. Collagen is the main structural protein in the ECM and connective tissue of animals. In mammals, collagen protein is highly abundant and mainly localized in the ECM of fibrous connective tissues, such as the tendon and skin [7–10]. It plays key structural roles by supporting the formation, tensile strength, and flexibility of joints [11–15]. Collagen types I, II, III, V, and XI are able to form fibrils that are necessary for structural support and resistance to mechanical stress in connective tissues [16,17]. Type I collagen is the most abundant form and is mainly present in the tendons and skin [18–20].

Collagen has numerous biomedical applications ranging from wound healing, bone and tissue regeneration, and drug delivery (Figure 1) [21,22]. Its accessibility, flexibility, and biocompatibility make it a useful biomaterial in several fields [22–24]. Collagen is a trimeric molecule made up of three polypeptide alpha-chains, forming highly organized three-dimensional structures capable of resisting mechanical stress and

supporting the growth of cells [25,26]. Marine organisms such as fish, jellyfish, sponges, and other invertebrates harbor a significant source of collagen and are highly advantageous over other sources, as they are metabolically compatible, lack religious constraints and are free of animal pathogens [27–30]. In fact, fish skins are commonly used for type I collagen extraction, as they are not only immensely abundant but also do not have religious restrictions and are not a risk of disease transmission [31–33]. Land animals possess many transmittable diseases, which makes them less favorable for use in industries. For example, cattle, although a large source of collagen, pose risks for bovine spongiform encephalopathy (BSE) as well as transmissible spongiform encephalopathy (TSE) [29,34,35]. These progressive neurological disorders affect cattle and can result in life-threatening infections in humans [29]. In addition, some religious constraints on the use of bovines for the pharmaceutical and cosmetic industries are up for debate [35]. These factors make marine sources of collagen a much safer, easier, and promising alternative. Marine organisms such as fish, jellyfish, sponges, and other invertebrates harbor significant source of collagen and are highly advantageous over other sources, as they metabolically compatible, lack religious constraints and are free of animal pathogens [27–30].

In fact, fish skins are commonly used for type I collagen extraction, as they are not only immensely abundant but also do not have religious restrictions and are not a risk disease transmission [31–33]. Land animals possess many transmittable diseases, which makes them less favorable for use in industries. For example, cattle, although a large source of collagen, pose risks for bovine spongiform encephalopathy (BSE) as well transmissible spongiform encephalopathy (TSE) [29,34,35]. These progressive neurological disorders affect cattle and can result in life-threatening infections in humans [29]. addition, some religious constraints on the use of bovines for the pharmaceutical and cosmetic industries are up for debate [35]. These factors make marine sources of collagen much safer, easier, and promising alternative. Skin wounds may take a long time to heal and often do not heal completely. Marine collagen isolated from organisms like fish, jellyfish, and sponges has been implicated several studies on its potential for increasing wound healing rates [36–41]. The processes involve increased fibroblast and keratinocyte migration as well as vascularization and growth [42–44]. In addition to accelerating wound healing, marine collagen has also been shown to have anti-aging properties by slowing the aging process in mice [45–48].

Studies on humans have also shown that marine collagen can reduce wrinkles, improve skin elasticity, and enhance the overall structure and appearance of skin. Furthermore, collagen's ability to regenerate bone has been shown to be successful in rat models of menopausal osteoporosis [49]. Marine collagen is able to increase bone mineral density and osteoblastic activity, serving protective effects against bone degeneration [49–53]. Collagen has also been shown to induce chondrogenic differentiation and prevent the development.Skin wounds may take a long time to heal and often do not heal completely. Marine collagen isolated from organisms like fish, jellyfish, and sponges has been implicated in several studies on its potential for increasing wound healing rates [36–41]. The processes involve increased fibroblast and keratinocyte migration as well as vascularization and epidermal growth [42–44]. In addition to accelerating wound healing, marine collagen has also been shown to have anti-aging properties by slowing the aging process in mice [45–48]. Studies on humans have also shown that marine collagen can reduce wrinkles, improve skin elasticity, and enhance the overall structure and appearance of skin. Furthermore, collagen's ability to regenerate bone has been shown to be successful in rat models of menopausal osteoporosis [49]. Marine collagen is able to increase bone mineral density and osteoblastic activity, serving protective effects against bone degeneration [49–53]. Collagen has also been shown to induce chondrogenic differentiation and prevent the development of osteoarthritis (OA) [54,55].



Technical advancements

Liposomal Technology: Vitamin C and Glutathione
Pepzyme Pro boosted Marine Collagen with Peach Extract,
Cold Water Soluble Vitamins

Composition of collafibe sachets

Nutritional Facts

Serving Size: 12g **Serving Per Container : 1**

		% ICMR RDA
Energy	31.0 Kcal	*
Total Carbohydrates	–	*
Total Sugar Includes 0g of added sugars	–	*
Protein	4.95 g	*
Sodium	20.1 mg	*
Fish Collagen Peptide Type 1	5.5 g	*
Lycopene 10% powder	5 mg	*
Grape seed extract 95%	10 mg	*
Peach extract	20 mg	*
Betaine Anhydrous (as Trimethylglycine)	20 mg	*
Niacinamide	20 mg	50%
Taurine	50 mg	*
Liposomal Vitamin C (as ascoribic acid) 70%	40 mg	50%
Vitamin E 50 CWS (as Tocopheryl Acetate) 50%	8 mg	80%
Sodium (as Sodium Chloride)	1.2 mg	0.06%
L-Lysine	5 mg	*
Beta carotene 10% Powder	1.2 mg	0.001%
Biotin	25 mcg	83.33%
Selenium (as L- Selenomethionine)	40 mcg	100%
Liposomal Glutathione 40%	250 mg	*

%ICMR RDA are Based on a 2000 calories diet.
***Indian Council of Medical Research Recommended Dietary Allowances Not Established**

Collafibe helps to:
Promote skin elasticity.
Promote skin hydration.
Reduce stretch marks.
Reduce visible cellulite.
Reduce dark spots.
Minimizing wrinkles and fine lines.
Protect from sun exposure.
Improve cell turnover and collagen formation, which keeps your skin stay elastic and healthy.

Collafibe sachets in wound healing and anti-aging

Our skin epidermis is the most important innate defense barrier against all pathogens and plays a significant role in tissue homeostasis [56–58]. Skin injuries are difficult to treat yet are becoming increasingly common as a result of burns, infections, scarring, genetic disorders, and other diseases [59,60]. Treatments aim to restore the integrity of the tissue, involving processes such as inflammation, cell division, differentiation, and vascularization. Endothelial permeability enables cell adhesion, which is followed by cell differentiation and maturation [61,62]. Marine collagen has been shown to be an effective biomaterial for wound healing. Collagen can be utilized in various formulations, such as the use of collagen peptides and hydroxylates, or collagen fibers, and scaffold-like structures [44,63]. Marine collagen peptides are produced from collagen through both chemical and enzymatic hydrolysis, and their smaller molecular weight increases their water solubility, making them more absorbable [63,64]. Hu et al. used an in vitro scratch assay to demonstrate that marine collagen peptides improve wound closure at concentrations of 50 _g mL□1 starting at 12 h post-treatment with collagen [63].

Collafibe sachets in wound healing and anti-aging

Our skin epidermis is the most important innate defense barrier against all pathogens and plays a significant role in tissue homeostasis [56–58]. Skin injuries are difficult to treat yet are becoming increasingly common as a result of burns, infections, scarring, genetic disorders, and other diseases [59,60]. Treatments aim to restore the integrity of the tissue, involving processes such as inflammation, cell division, differentiation, and vascularization. Endothelial permeability enables cell adhesion, which is followed by cell differentiation and maturation [61,62]. Marine collagen has been shown to be an effective biomaterial for wound healing. Collagen can be utilized in various formulations, such as the use of collagen peptides and hydroxylates, or collagen fibers, and scaffold-like structures [44,63]. Marine collagen peptides are produced from collagen through both chemical and enzymatic hydrolysis, and their smaller molecular weight increases their water solubility, making them more absorbable [63,64]. Hu et al. used an in vitro scratch assay to demonstrate that marine collagen peptides improve

wound closure at concentrations of 50 μg mL−1 starting at 12 h post-treatment with collagen [63]. Wang et al. found that marine collagen peptides (MCPs) isolated from salmon skin significantly improved skin wound tensile strength in rats [42].

The potential role of collafibe sachets in bone and cartilage regeneration

Marine collagen sources serve not only as a promising avenue for healing skin injuries but also for bone-related trauma and regeneration. Bone fracture repair and healing is a form of tissue regeneration and is a complex process involving bone formation and breakdown [90,91]. Often, patients present with conditions that require reconstruction of large bones as a result of genetic abnormalities, trauma, infection, and tumors [92]. There is an increasing demand to improve methods of bone repair and regeneration, such as functional bone grafts [93]. Marine collagen bioactive peptides are known to aid in the absorption of calcium and zinc, which are important components of bone and are beneficial for osteoporosis prevention [94,95]. A study performed by Xu et al found that marine collagen peptides isolated and derived by hydrolysis from chum salmon increased serum osteocalcin in treated rats compared to controls. Osteocalcin is a protein hormone secreted by osteoblasts and plays a role in bone maintenance and regeneration through interaction with calcium. The study also found that bone organic matrix, density, femoral length, and femur mineral ions were significantly higher in the collagen-treated group than in the controls [94].

The potential role of collagen in collafibe sachets in bone and cartilage regeneration

Marine collagen sources serve not only as a promising avenue for healing skin injuries but also for bone-related trauma and regeneration. Bone fracture repair and healing is a form of tissue regeneration and is a complex process involving bone formation and breakdown [90,91]. Often, patients present with conditions that require reconstruction of large bones as a result of genetic abnormalities, trauma, infection, and tumors [92]. There is an increasing demand to improve methods of bone repair and regeneration, such as functional bone grafts [93]. Marine collagen bioactive peptides are known to aid in the absorption of calcium and zinc, which are important components of bone and are beneficial for osteoporosis prevention [94,95]. A study performed by Xu et al found that marine collagen peptides isolated and derived by hydrolysis from chum salmon increased serum osteocalcin in treated rats compared to controls. Osteocalcin is a protein hormone secreted by osteoblasts and plays a role in bone maintenance and regeneration through interaction with calcium. The study also found that bone organic matrix, density, femoral length, and femur mineral ions were significantly higher in the collagen-treated group than in the controls [94]. It was hypothesized that the increase in bone mineral density was likely due to increased osteoblast activity, as seen by the increase in bone size and serum osteocalcin [94]. These results shed light on the potential collagen peptides involved in mineral deposition, bone matrix development and an increase in osteoblastic activity, which strongly suggests that collagen is a promising biomaterial for the prevention and treatment of osteoporosis [94]. Osteoporosis and net bone loss are prevalent among aging women going through menopause resulting from estrogen deficiency [49]. Nomura et al. demonstrated that 20 mg of collagen isolated from shark gelatin also increased the bone mineral density of the spongy bone in rat models of menopausal osteoporosis [49]. Furthermore, the biological effect of marine collagen on rat-derived bone marrow stem cells has also been demonstrated. Liu et al. showed that 0.2 mg/mL collagen isolated from fish promoted cell survival and upregulated the expression of several osteogenic and endothelial markers [50].

CONCLUSION

Studies on humans have also shown that marine collagen in Collafibe sachets can reduce wrinkles, improve skin elasticity, and enhance the overall structure and appearance of skin. Furthermore, collagen's ability to regenerate bone has been shown to be successful in rat models of menopausal osteoporosis. Marine collagen in Collafibe sachets is able to increase bone mineral density and osteoblastic activity, serving protective effects against bone degeneration. Collagen has also been shown to induce chondrogenic differentiation and prevent the development. Skin wounds may take a long time to heal and often do not heal completely. Marine collagen isolated from organisms like fish, jellyfish, and sponges has been implicated in several studies on its potential for increasing wound healing rates. The processes involve increased fibroblast and keratinocyte migration as well as vascularization and epidermal growth. In addition to accelerating wound healing, marine collagen has also been shown to have anti-aging properties by slowing the aging process.

REFERENCES

1. Rahman, M.A. Collagen of Extracellular Matrix from Marine Invertebrates and Its Medical Applications. Mar. Drugs 2019, 17, 118.
2. Rodriguez-Pascual, F.; Slatter, D.A. Collagen Cross-Linking: Insights on the Evolution of Metazoan Extracellular Matrix. Sci. Rep. 2016, 6, 37374.

3. Frantz, C.; Stewart, K.M; Weaver, V.M. The Extracellular Matrix at a Glance. J. Cell Sci. 2010, 123, 4195–4200.
4. Zhang, Y.; He, Y.; Bharadwaj, S.; Hammam, N.; Carnagey, K.; Myers, R.; Atala, A.; Van Dyke, M. Tissue-Specific Extracellular Matrix Coatings for the Promotion of Cell Proliferation and Maintenance of Cell Phenotype. Biomaterials 2009, 30, 4021–4028.
5. Gérard, C.; Goldbeter, A. The Balance between Cell Cycle Arrest and Cell Proliferation: Control by the Extracellular Matrix and by Contact Inhibition. Interface Focus 2014, 4, 20130075.
6. Townsend, S.E.; Gannon, M. Extracellular Matrix–Associated Factors Play Critical Roles in Regulating Pancreatic _-Cell Proliferation and Survival. Endocrinology 2019, 160, 1885–1894.
7. Hahn, M.S.; Kobler, J.B.; Zeitels, S.M.; Langer, R. Quantitative and Comparative Studies of the Vocal Fold Extracellular Matrix II: Collagen. Ann. Otol. Rhinol. Laryngol. 2006, 115, 225–232.
8. Birch, H.L. Extracellular Matrix and Ageing. In Biochemistry and Cell Biology of Ageing: Part I Biomedical Science; Harris, J.R., Korolchuk, V.I., Eds.; Subcellular Biochemistry; Springer: Singapore, 2018; Volume 90, pp. 169–190, ISBN 9789811328343.
9. Gelse, K. Collagens—Structure, Function, and Biosynthesis. Adv. Drug Deliv. Rev. 2003, 55, 1531–1546.
10. Sorushanova, A.; Delgado, L.M.; Wu, Z.; Shologu, N.; Kshirsagar, A.; Raghunath, R.; Mullen, A.M.; Bayon, Y.; Pandit, A.;Raghunath, M.; et al. The Collagen Suprafamily: From Biosynthesis to Advanced Biomaterial Development. Adv. Mater. 2019,31, 1801651.
11. Whelan, A.; Duffy, J.; Gaul, R.T.; O'Reilly, D.; Nolan, D.R.; Gunning, P.; Lally, C.; Murphy, B.P. Collagen Fibre Orientation and Dispersion Govern Ultimate Tensile Strength, Stiffness and the Fatigue Performance of Bovine Pericardium. J. Mech. Behav. Biomed. Mater. 2019, 90, 54–60.
12. Miguez, P.A.; Pereira, P.N.R.; Atsawasuwan, P.; Yamauchi, M. Collagen Cross-Linking and Ultimate Tensile Strength in Dentin. J. Dent. Res. 2004, 83, 807–810.
13. Raz, P.; Brosh, T.; Ronen, G.; Tal, H. Tensile Properties of Three Selected Collagen Membranes. BioMed Res. Int. 2019, 2019,5163603.
14. Clark, K.L.; Sebastianelli,W.; Flechsenhar, K.R.; Aukermann, D.F.; Meza, F.; Millard, R.L.; Deitch, J.R.; Sherbondy, P.S.; Albert, A. 24-Week Study on the Use of Collagen Hydrolysate as a Dietary Supplement in Athletes with Activity-Related Joint Pain. Curr. Med. Res. Opin. 2008, 24, 1485–1496.
15. Zdzieblik, D.; Oesser, S.; Gollhofer, A.; König, D. Improvement of Activity-Related Knee Joint Discomfort Following Supplementationof Specific Collagen Peptides. Appl. Physiol. Nutr. Metab. 2017, 42, 588–595.
16. Holmes, D.F.; Lu, Y.; Starborg, T.; Kadler, K.E. Collagen Fibril Assembly and Function. In Current Topics in Developmental Biology;Elsevier: Amsterdam, The Netherlands, 2018; Volume 130, pp. 107–142, ISBN 978-0-12-809802-8.
17. Wess, T.J. Collagen Fibril Form and Function. In Advances in Protein Chemistry; Elsevier: Amsterdam, The Netherlands, 2005; Volume 70, pp. 341–374, ISBN 978-0-12-034270-9.Mar. Drugs 2022, 20, 61 13 of 16
18. Chowdhury, S.R.; Mh Busra, M.F.; Lokanathan, Y.; Ng, M.H.; Law, J.X.; Cletus, U.C.; Binti Haji Idrus, R. Collagen Type I: A Versatile Biomaterial. In Novel Biomaterials for Regenerative Medicine; Chun, H.J., Park, K., Kim, C.-H., Khang, G., Eds.; Advances in Experimental Medicine and Biology; Springer: Singapore, 2018; Volume 1077, pp. 389–414, ISBN 9789811309465.
19. Kisling, A.; Lust, R.M.; Katwa, L.C. What Is the Role of Peptide Fragments of Collagen I and IV in Health and Disease? Life Sci.2019, 228, 30–34.
20. Nicol, L.; Morar, P.;Wang, Y.; Henriksen, K.; Sun, S.; Karsdal, M.; Smith, R.; Nagamani, S.C.S.; Shapiro, J.; Lee, B.; et al. Alterations in Non-Type I Collagen Biomarkers in Osteogenesis Imperfecta. Bone. 2019, 120, 70–74.
21. Ferrario, C.; Rusconi, F.; Pulaj, A.; Macchi, R.; Landini, P.; Paroni, M.; Colombo, G.; Martinello, T.; Melotti, L.; Gomiero, C.; et al. From Food Waste to Innovative Biomaterial: Sea Urchin-Derived Collagen for Applications in Skin Regenerative Medicine. Mar. Drugs 2020, 18, 414.
22. Subhan, F.; Ikram, M.; Shehzad, A.; Ghafoor, A. Marine Collagen: An Emerging Player in Biomedical Applications. J. Food Sci.Technol. 2015, 52, 4703–4707.
23. Benayahu, D.; Pomeraniec, L.; Shemesh, S.; Heller, S.; Rosenthal, Y.; Rath-Wolfson, L.; Benayahu, Y. Biocompatibility of a MarineCollagen-Based Scaffold In Vitro and In Vivo. Mar. Drugs 2020, 18, 420.
24. Song, E.; Yeon Kim, S.; Chun, T.; Byun, H.-J.; Lee, Y.M. Collagen Scaffolds Derived from a Marine Source and Their Biocompatibility.Biomaterials 2006, 27, 2951–2961.
25. Boudko, S.P.; Sasaki, T.; Engel, J.; Lerch, T.F.; Nix, J.; Chapman, M.S.; Bächinger, H.P. Crystal Structure of Human Collagen XVIII Trimerization Domain: A Novel Collagen Trimerization Fold. J. Mol. Biol. 2009, 392, 787–802. [PubMed]
26. Boudko, S.P.; Engel, J.; Bächinger, H.P. The Crucial Role of Trimerization Domains in Collagen Folding. Int. J. Biochem. Cell Biol. 2012, 44, 21–32.

27. Sugiura, H.; Yunoki, S.; Kondo, E.; Ikoma, T.; Tanaka, J.; Yasuda, K. In Vivo Biological Responses and Bioresorption of Tilapia Scale Collagen as a Potential Biomaterial. J. Biomater. Sci. Polym. Ed. 2009, 20, 1353–1368.
28. Cheng, X.; Shao, Z.; Li, C.; Yu, L.; Raja, M.A.; Liu, C. Isolation, Characterization and Evaluation of Collagen from JellyfishRhopilema Esculentum Kishinouye for Use in Hemostatic Applications. PLoS ONE 2017, 12, e0169731.
29. Widdowson, J.P.; Picton, A.J.; Vince, V.; Wright, C.J.; Mearns-Spragg, A. In Vivo Comparison of Jellyfish and Bovine Collagen Sponges as Prototype Medical Devices: In Vivo comparison of jellyfish and bovine collagen sponges. J. Biomed. Mater. Res. B Appl. Biomater. 2018, 106, 1524–1533. [PubMed]
30. Coppola, D.; Oliviero, M.; Vitale, G.A.; Lauritano, C.; D'Ambra, I.; Iannace, S.; de Pascale, D. Marine Collagen from Alternative and Sustainable Sources: Extraction, Processing and Applications. Mar. Drugs 2020, 18, 214. [PubMed]
31. Silvipriya, K.; Kumar, K.; Bhat, A.; Kumar, B.; John, A.; Lakshmanan, P. Collagen: Animal Sources and Biomedical Application. J. Appl. Pharm. Sci. 2015, 5, 123–127.
32. Shavandi, A.; Hou, Y.; Carne, A.; McConnell, M.; Bekhit, A.E.-D.A. Marine Waste Utilization as a Source of Functional and Health Compounds. Adv. Food Nutr. Res. 2019, 87, 187–254.
33. Hou, Y.; Shavandi, A.; Carne, A.; Bekhit, A.A.; Ng, T.B.; Cheung, R.C.F.; Bekhit, A.E.A. Marine Shells: Potential Opportunities for Extraction of Functional and Health-Promoting Materials. Crit. Rev. Environ. Sci. Technol. 2016, 46, 1047–1116.
34. Karim, A.A.; Bhat, R. Gelatin Alternatives for the Food Industry: Recent Developments, Challenges and Prospects. Trends FoodSci. Technol. 2008, 19, 644–656.
35. Karim, A.A.; Bhat, R. Fish Gelatin: Properties, Challenges, and Prospects as an Alternative to Mammalian Gelatins. Food Hydrocoll. 2009, 23, 563–576.
36. Yamada, S.; Yamamoto, K.; Ikeda, T.; Yanagiguchi, K.; Hayashi, Y. Potency of Fish Collagen as a Scaffold for Regenerative Medicine. BioMed Res. Int. 2014, 2014, 302932.
37. Zhou, T.;Wang, N.; Xue, Y.; Ding, T.; Liu, X.; Mo, X.; Sun, J. Electrospun Tilapia Collagen Nanofibers AcceleratingWound Healing via Inducing Keratinocytes Proliferation and Differentiation. Colloids Surf. B Biointerfaces 2016, 143, 415–422. [PubMed]
38. Sampath Kumar, N.S.; Nazeer, R.A.; Jaiganesh, R.Wound Healing Properties of Collagen from the Bone of Two Marine Fishes. Int. J. Pept. Res. Ther. 2012, 18, 185–192.
39. Shalaby, M.; Agwa, M.; Saeed, H.; Khedr, S.M.; Morsy, O.; El-Demellawy, M.A. Fish Scale Collagen Preparation, Characterization and Its Application in Wound Healing. J. Polym. Environ. 2020,28,166-78.
40. Feng, X.; Zhang, X.; Li, S.; Zheng, Y.; Shi, X.; Li, F.; Guo, S.; Yang, J. Preparation of Aminated Fish Scale Collagen and Oxidized Sodium Alginate Hybrid Hydrogel for Enhanced Full-Thickness Wound Healing. Int. J. Biol. Macromol. 2020, 164, 626–637. [PubMed]
41. Pal, P.; Srivas, P.K.; Dadhich, P.; Das, B.; Maity, P.P.; Moulik, D.; Dhara, S. Accelerating Full Thickness Wound Healing Using Collagen Sponge of Mrigal Fish (Cirrhinus Cirrhosus) Scale Origin. Int. J. Biol. Macromol. 2016, 93, 1507–1518. [PubMed]
42. Wang, J.; Xu, M.; Liang, R.; Zhao, M.; Zhang, Z.; Li, Y. Oral Administration of Marine Collagen Peptides Prepared from Chum Salmon (Oncorhynchus Keta) Improves Wound Healing Following Cesarean Section in Rats. Food Nutr. Res. 2015, 59, 26411.
43. Zhang, Z.; Wang, J.; Ding, Y.; Dai, X.; Li, Y. Oral Administration of Marine Collagen Peptides from Chum Salmon Skin Enhances Cutaneous Wound Healing and Angiogenesis in Rats. J. Sci. Food Agric. 2011, 91, 2173–2179. Mar. Drugs 2022, 20, 61 14 of 16
44. Chen, J.; Gao, K.; Liu, S.; Wang, et al. Fish Collagen Surgical Compress Repairing Characteristics on Wound Healing Process In Vivo. Mar. Drugs 2019, 17, 33.
45. Ito, N.; Seki, S.; Ueda, F. Effects of Composite Supplement Containing Collagen Peptide and Ornithine on Skin Conditions and Plasma IGF-1 Levels—A Randomized, Double-Blind, Placebo-Controlled Trial. Mar. Drugs 2018, 16, 482.
46. De Luca, C.; Mikhal'chik, E.V.; Suprun, M.V.; Papacharalambous, M.; Truhanov, A.I.; Korkina, L.G. Skin Antiageing and Systemic Redox Effects of Supplementation with Marine Collagen Peptides and Plant-Derived Antioxidants: A Single-Blind Case-Control Clinical Study. Oxid. Med. Cell. Longev. 2016, 2016, 4389410. [PubMed]
47. Allouche, M.; Hamdi, I.; Nasri, A.; Harrath, A.H.; Mansour, L.; Beyrem, H.; Boufahja, F. Laboratory Bioassay Exploring the Effects of Anti-Aging Skincare Products on Free-Living Marine Nematodes: A Case Study of Collagen. Environ. Sci. Pollut. Res. 2020, 27, 11403–11412.
48. Kim, C.-R.; Kim, Y.-M.; Lee, M.-K.; Kim, I.-H.; Choi, Y.-H.; Nam, T.-J. Pyropia Yezoensis Peptide Promotes Collagen Synthesis by Activating the TGF-_/Smad Signaling Pathway in the Human Dermal Fibroblast Cell Line Hs27. Int. J. Mol. Med. 2017, 39, 31–38.

49. Nomura, Y.; Oohashi, K.; Watanabe, M.; Kasugai, S. Increase in Bone Mineral Density through Oral Administration of Shark Gelatin to Ovariectomized Rats. Nutrition 2005, 21, 1120–1126.
50. Liu, C.; Sun, J. Potential Application of Hydrolyzed Fish Collagen for Inducing the Multidirectional Differentiation of Rat Bone Marrow Mesenchymal Stem Cells. Biomacromolecules 2014, 15, 436–443.
51. Yamada, S.; Nagaoka, H.; Terajima, M.; Tsuda, N.; Hayashi, Y.; Yamauchi, M. Effects of Fish Collagen Peptides on Collagen Post-Translational Modifications and Mineralization in an Osteoblastic Cell Culture System. Dent. Mater. J. 2013, 32, 88–95.
52. Elango, J.; Robinson, J.; Zhang, J.; Bao, B.; Ma, N.; de Val, J.E.M.S.; Wu, W. Collagen Peptide Upregulates Osteoblastogenesis from Bone Marrow Mesenchymal Stem Cells through MAPK- Runx2. Cells 2019, 8, 446.
53. Capati, M.L.F.; Nakazono, A.; Yamamoto, K.; Sugimoto, K.; Yanagiguchi, K.; Yamada, S.; Hayashi, Y. Fish Collagen Promotes the Expression of Genes Related to Osteoblastic Activity. Int. J. Polym. Sci. 2016, 2016, 5785819.
54. Hsu, H.-H.; Uemura, T.; Yamaguchi, I.; Ikoma, T.; Tanaka, J. Chondrogenic Differentiation of Human Mesenchymal Stem Cells on Fish Scale Collagen. J. Biosci. Bioeng. 2016, 122, 219–225.
55. Bourdon, B.; Contentin, R.; Cassé, F.; Maspimby, C.; Oddoux, S.; Noël, A.; Legendre, F.; Gruchy, N.; Galéra, P. Marine Collagen Hydrolysates Downregulate the Synthesis of Pro-Catabolic and Pro-Inflammatory Markers of Osteoarthritis and Favor Collagen Production and Metabolic Activity in Equine Articular Chondrocyte Organoids. Int. J. Mol. Sci. 2021, 22, 580.
56. Blanpain, C.; Fuchs, E. Epidermal Homeostasis: A Balancing Act of Stem Cells in the Skin. Nat. Rev. Mol. Cell Biol. 2009, 10, 207–217. [PubMed]
57. Elias, P.M. The Skin Barrier as an Innate Immune Element. Semin. Immunopathol. 2007, 29, 3–14.
58. Harder, J.; Schröder, J.-M.; Gläser, R. The Skin Surface as Antimicrobial Barrier: Present Concepts and Future Outlooks. Exp. Dermatol. 2013, 22, 1–5.
59. Shakespeare, P.G. The Role of Skin Substitutes in the Treatment of Burn Injuries. Clin. Dermatol. 2005, 23, 413–418.
60. Hall, A.H.; Mathieu, L.; Maibach, H.I. Acute Chemical Skin Injuries in the United States: A Review. Crit. Rev. Toxicol. 2018, 48, 540–554. [PubMed]
61. Cerutti, C.; Ridley, A.J. Endothelial Cell-Cell Adhesion and Signaling. Exp. Cell Res. 2017, 358, 31–38.
62. Han, J.; Shuvaev, V.V.; Muzykantov, V.R. Catalase and Superoxide Dismutase Conjugated with Platelet-Endothelial Cell Adhesion Molecule Antibody Distinctly Alleviate Abnormal Endothelial Permeability Caused by Exogenous Reactive Oxygen Species and Vascular Endothelial Growth Factor. J. Pharmacol. Exp. Ther. 2011, 338, 82–91. [PubMed]
63. Hu, Z.; Yang, P.; Zhou, C.; Li, S.; Hong, P. Marine Collagen Peptides from the Skin of Nile Tilapia (Oreochromis Niloticus):Characterization and Wound Healing Evaluation. Mar. Drugs 2017, 15, 102.
64. León-López, A.; Morales-Peñaloza, A.; Martínez-Juárez, V.M.; Vargas-Torres, A.; Zeugolis, D.I.; Aguirre-Álvarez, G. Hydrolyzed Collagen—Sources and Applications. Molecules 2019, 24, 4031.
65. Yang, T.; Zhang, K.; Li, B.; Hou, H. Effects of Oral Administration of Peptides with Low MolecularWeight from Alaska Pollock (Theragra Chalcogramma) on Cutaneous Wound Healing. J. Funct. Foods 2018, 48, 682–691.
66. Pozzolini, M.; Millo, E.; Oliveri, C.; Mirata, S.; Salis, A.; Damonte, G.; Arkel, M.; Scarfì, S. Elicited ROS Scavenging Activity, Photoprotective, and Wound-Healing Properties of Collagen-Derived Peptides from the Marine Sponge Chondrosia Reniformis. Mar. Drugs 2018, 16, 465.
67. Felician, F.F.; Yu, R.-H.; Li, M.-Z.; Li, C.-J.; Chen, H.-Q.; Jiang, Y.; Tang, T.; Qi,W.-Y.; Xu, H.-M. TheWound Healing Potential of Collagen Peptides Derived from the Jellyfish Rhopilema Esculentum. Chin. J. Traumatol. 2019, 22, 12–20. [PubMed]
68. Veeruraj, A.; Liu, L.; Zheng, J.; Wu, J.; Arumugam, M. Evaluation of Astaxanthin Incorporated Collagen Film Developed from the Outer Skin Waste of Squid Doryteuthis Singhalensis for Wound Healing and Tissue Regenerative Applications. Mater. Sci. Eng. C 2019, 95, 29–42.
69. Melotti, L.; Martinello, T.; Perazzi, A.; Iacopetti, I.; Ferrario, C.; Sugni, M.; Sacchetto, R.; Patruno, M. A Prototype Skin Substitute, Made of Recycled Marine Collagen, Improves the Skin Regeneration of Sheep. Animals 2021, 11, 1219.
70. Farage, M.A.; Miller, K.W.; Elsner, P.; Maibach, H.I. Characteristics of the Aging Skin. Adv. Wound Care 2013, 2, 5–10. Mar. Drugs 2022, 20, 61 15 of 16
71. Oxlund, H.; Manschot, J.; Viidik, A. The Role of Elastin in the Mechanical Properties of Skin. J. Biomech. 1988, 21, 213–218.
72. Aziz, J.; Shezali, H.; Radzi, Z.; Yahya, N.A.; Abu Kassim, N.H.; Czernuszka, J.; Rahman, M.T. Molecular Mechanisms of
73. Stress-Responsive Changes in Collagen and Elastin Networks in Skin. Skin Pharmacol. Physiol. 2016, 29, 190–203.

74. Varani, J.; Dame, M.K.; Rittie, L.; Fligiel, S.E.G.; Kang, S.; Fisher, G.J.; Voorhees, J.J. Decreased Collagen Production in Chronologically Aged Skin. Am. J. Pathol. 2006, 168, 1861–1868. [PubMed]
75. Mays, P.; Bishop, J.; Laurent, G. Age-Related Changes in the Proportion of Types I and III Collagen. Mech. Ageing Dev. 1988, 45, 203–212.
76. Boelsma, E.; Hendriks, H.F.; Roza, L. Nutritional Skin Care: Health Effects of Micronutrients and Fatty Acids. Am. J. Clin. Nutr. 2001, 73, 853–864.
77. Pullar, J.; Carr, A.; Vissers, M. The Roles of Vitamin C in Skin Health. Nutrients 2017, 9, 866.
78. Ben-Shabat, S.; Kazdan, Y.; Beit-Yannai, E.; Sintov, A.C. Use of Alpha-Tocopherol Esters for Topical Vitamin E Treatment: Evaluation of Their Skin Permeation and Metabolism. J. Pharm. Pharmacol. 2013, 65, 652–658.
79. Vaiserman, A.; Koliada, A.; Zayachkivska, A.; Lushchak, O. Nanodelivery of Natural Antioxidants: An Anti-Aging Perspective. Front. Bioeng. Biotechnol. 2020, 7, 447. [PubMed]
80. Luo, J.; Zhou, Z.; Yao, X.; Fu, Y. Mineral-Chelating Peptides Derived from Fish Collagen: Preparation, Bioactivity and Bioavailability. LWT 2020, 134, 110209.
81. Exposito, J.-Y.; Valcourt, U.; Cluzel, C.; Lethias, C. The Fibrillar Collagen Family. Int. J. Mol. Sci. 2010, 11, 407–426.
82. Alves, A.; Marques, A.; Martins, E.; Silva, T.; Reis, R. Cosmetic Potential of Marine Fish Skin Collagen. Cosmetics 2017, 4, 39.
83. Evans, M.; Lewis, E.D.; Zakaria, N.; Pelipyagina, T.; Guthrie, N. A Randomized, Triple-blind, Placebo-controlled, Parallel Study to Evaluate the Efficacy of a Freshwater Marine Collagen on SkinWrinkles and Elasticity. J. Cosmet. Dermatol. 2021, 20, 825–834.
84. Wang, Z.;Wang, Q.;Wang, L.; Xu,W.; He, Y.; Li, Y.; He, S.; Ma, H. Improvement of Skin Condition by Oral Administration of Collagen Hydrolysates in Chronologically Aged Mice: Skin Improvement by Collagen Hydrolysates in Chronologically Aged Mice. J. Sci. Food Agric. 2017, 97, 2721–2726.
85. Pei, X.; Yang, R.; Zhang, Z.; Xu, Y.; Han, X.;Wang, J.; Li, Y. Effects of marine collagen peptide on delaying the skin aging. Zhonghua Yu Fang Yi Xue Za Zhi [Chin. J. Prev. Med.] 2008, 42, 235–238.
86. Davies, K.J.A.; Quintanilha, A.T.; Brooks, G.A.; Packer, L. Free Radicals and Tissue Damage Produced by Exercise. Biochem. Biophys. Res. Commun. 1982, 107, 1198–1205.
87. Sailaja Rao, P.; Kalva, S.; Yerramilli, A.; Mamidi, S. Free Radicals and Tissue Damage: Role of Antioxidants. Free Radic. Antioxid. 2011, 1, 2–7.
88. Rinnerthaler, M.; Bischof, J.; Streubel, M.; Trost, A.; Richter, K. Oxidative Stress in Aging Human Skin. Biomolecules 2015, 5,545–589.
89. Venkatesan, J.; Anil, S.; Kim, S.-K.; Shim, M. Marine Fish Proteins and Peptides for Cosmeceuticals: A Review. Mar. Drugs 2017,15, 143. [PubMed]
90. Chi, C.-F.; Hu, F.-Y.; Wang, B.; Ren, X.-J.; Deng, S.-G.; Wu, C.-W. Purification and Characterization of Three Antioxidant Peptides from Protein Hydrolyzate of Croceine Croaker (Pseudosciaena Crocea) Muscle. Food Chem. 2015, 168, 662–667.
91. Hadjiargyrou, M.; Lombardo, F.; Zhao, S.; Ahrens, W.; Joo, J.; Ahn, H.; Jurman, M.; White, D.W.; Rubin, C.T. Transcriptional Profiling of Bone Regeneration. J. Biol. Chem. 2002, 277, 30177–30182.
92. Armiento, A.R.; Hatt, L.P.; Sanchez Rosenberg, G.; Thompson, K.; Stoddart, M.J. Functional Biomaterials for Bone Regeneration: A Lesson in Complex Biology. Adv. Funct. Mater. 2020, 30, 1909874.
93. Dimitriou, R.; Jones, E.; McGonagle, D.; Giannoudis, P.V. Bone Regeneration: Current Concepts and Future Directions. BMC Med.2011, 9, 66.
94. Cicciù, M. Real Opportunity for the Present and a Forward Step for the Future of Bone Tissue Engineering. J. Craniofac. Surg. 2017, 28, 592–593.
95. Xu, Y.; Han, X.; Li, Y. Effect of Marine Collagen Peptides on Long Bone Development in Growing Rats. J. Sci. Food Agric. 2010, 90, 1485–1491.
96. Meng, K.; Chen, L.; Xia, G.; Shen, X. Effects of Zinc Sulfate and Zinc Lactate on the Properties of Tilapia (Oreochromis Niloticus) Skin Collagen Peptide Chelate Zinc. Food Chem. 2021, 347, 129043.